

## Screening Level Ecological Risk Assessment — Final

**Prepared for:**

Utah Department of Environmental Quality  
Division of Water Quality

**Chevron Pipeline June 2010 Crude Oil Spill  
Lower Red Butte Creek  
Salt Lake City, Utah**

November 2012

[www.erm.com](http://www.erm.com)



Utah Department of Environmental Quality  
Division of Water Quality

## Screening Level Ecological Risk Assessment — Final

*Chevron Pipeline June 2010 Crude Oil Spill  
Lower Red Butte Creek  
Salt Lake City, Utah*

November 2012

Project No. 0145323

---

Brent Robinson  
*Partner-in-Charge*

---

Mark Shibata  
*Program/Technical Director*

---

Sandra Mulhearn  
*Senior Scientist*

**ERM-West, Inc.**  
102 West 500 South  
Salt Lake City, UT 84101-2334  
T: 801-595-8400  
F: 801-595-8484

• This page intentionally left blank •

## TABLE OF CONTENTS

Appendices .....	v
List of Tables .....	vi
List of Figures .....	vii
List of Aconyms and Abbreviations.....	viii
Executive Summary .....	xi
1.0 Introduction .....	1
1.1 Spill History .....	1
1.2 Overview of The Ecological Risk Assessment .....	2
2.0 Site Description .....	5
2.1 Location / General Setting.....	5
2.2 Geology .....	6
2.3 In-Creek Flows.....	6
2.4 Flora and Fauna.....	6
3.0 Data Evaluation.....	9
3.1 Data Collection .....	9
3.2 Data Validation.....	12
3.3 Data Usability .....	13
3.4 Handling Non-Detected Values.....	16
4.0 Local background Evaluation .....	19
4.1 Statistical Comparisons .....	20
4.2 Comparision to Background Threshold Values.....	21
4.3 polycyclic aromatic hydrocarbons Composition Analysis.....	22
4.4 Summary .....	23
5.0 Problem Formulation .....	25
5.1 Biotic Receptors of Concern.....	25
5.2 Constituents of Potential Ecological Concern.....	27
5.3 Potentially Complete Exposure Pathways .....	29
5.4 Assessment Endpoints.....	31

6.0	Exposure Assessment.....	33
6.1	Total Petroleum hydrocarbons Concentrations.....	33
6.2	Exposure Point Concentrations.....	33
6.3	Aquatic and Benthic Macroinvertebrates .....	34
6.4	Riparian Wildlife .....	34
7.0	Effects Assessment.....	39
7.1	Aquatic Biota toxicity reference values.....	40
7.2	Benthic Macroinvertebrate Toxicity Reference Values .....	40
7.3	Riparian Wildlife Toxicity Reference Values .....	41
7.4	In-Creek Benthic Macroinvertebrate Community Structure .....	42
8.0	Risk Characterization .....	45
8.1	Hazard Quotients .....	45
8.2	Aquatic Biota.....	46
8.3	Benthic Macroinvertebrates .....	46
8.4	Riparian Birds .....	52
8.5	Riparian Mammals.....	54
8.6	Uncertainty Analysis .....	57
9.0	Discussion and Conclusions.....	65
10.0	Lituration Cited.....	67

***APPENDICES***

- Appendix A: Ecological Risk Assessment Work Plan
- Appendix B: Photographs of Sampling Locations
- Appendix C: Quality Assurance/Quality Control Memorandums
- Appendix D: Local Background Evaluation
- Appendix E: Ecological Risk Analyses
- Appendix F: Analysis of Benthic Macroinvertebrate Community Indices
- Appendix G: Data Used in Risk Assessment (provided by McDaniel-Lambert) [see attached CD]
- Appendix H: Location-Specific Risk Estimates
- Appendix I: Responses to Public Comments

**LIST OF TABLES**

Table 3-1.	Spill Monitoring Sampling and Analysis Plan and Supplemental Sampling Stations to Support the Ecological Risk Assessment
Table 3-2.	Criteria Used to Identify Data Usability
Table 5-1.	Risk-Based Ecological Screening Levels
Table 5-2.	Constituents of Potential Ecological Concern
Table 5-3.	Assessment Endpoints for Lower (Urban) Red Butte Creek
Table 6-1.	Wildlife Exposure Factors
Table 6-2.	Bioaccumulation Factors
Table 7-1.	Sources of Toxicity Reference Values
Table 7-2.	Aquatic Biota Toxicity Reference Values
Table 7-3.	Benthic Macroinvertebrate Community Toxicity Reference Values
Table 7-4.	Wildlife Toxicity Reference Values
Table 7-5.	In-Creek Benthic Macroinvertebrate Community Metrics
Table 8-1.	Risk Calculations for Aquatic Biota
Table 8-2.	Risk Calculations for Benthic Macroinvertebrates
Table 8-3.	Risk Calculations for the Mallard
Table 8-4.	Risk Calculations for the Sandpiper
Table 8-5.	Risk Calculations for the Muskrat
Table 8-6.	Risk Calculations for the Raccoon
Table 8-7.	Benthic Macroinvertebrate Sampling Locations within Lower Red Butte Creek
Table 8-8.	Benthic Macroinvertebrate Community: Results of Nonparametric Two-Sample Tests
Table 8-9.	Benthic Macroinvertebrate Community: Results of Nonparametric Regression Analyses
Table 8-10.	Summary of Location-Specific Risk Estimates for TPH

**Note:** Tables 8-1 through 8-6 found at end of the document.



***LIST OF FIGURES***

Figure 1-1. ERA Approach

Figure 1-2. Elements of the SLERA

Figure 2-1. Map of Red Butte, Emigration, City, and Parleys Creeks

Figure 2-2. Longitudinal Profile Plot of Lower Red Butte Creek Streambed

Figure 3-1. Map of Sampling Locations

Figure 5-1. Conceptual Site Model for the Lower Red Butte Creek SLERA

Figure 8-1. Benthic Macroinvertebrate: Boxplots of Community Metrics

Figure 8-2. Map of Location-Specific Risk Estimates

**LIST OF ACRONYMS AND ABBREVIATIONS**

95UCL	95 percent upper confidence limits
AF	assimilation factor
ARCS	Assessment and Remediation Contaminated Sediments
ATSDR	Agency for Toxic Substances and Disease Registry
BAF	bioaccumulation factor
BERA	baseline ecological risk assessment
BROC	biotic receptor of concern
BTEX	benzene, toluene, ethylbenzene, xylene
BTV	background threshold value
BW	body weight
CD	compact disk
cfs	cubic feet per second
COPEC	constituent of potential ecological concern
CPL	Chevron Pipeline Company
CR	contact rate
CSM	conceptual site model
DQO	data quality objective
E <sub>s</sub>	sorting effectiveness
EPC	exposure point concentration
EPT	ephemeroptera, plecoptera, trichoptera
ERA	ecological risk assessment
ERM	ERM-West, Inc.
ESL	ecological screening level
ESSL	ecological soil screening level
FC	fraction of media contacted
H <sub>o</sub>	null hypothesis
ha	hectare
HBI	Hilsenhoff biotic index
HHRA	human health risk assessment
HI	hazard index
HQ	hazard quotient
kg	kilogram

L	liter
LD <sub>50</sub>	lethal dose for 50 percent of the population
LOAEL	lowest observable adverse effect level
MaDEP	Massachusetts Department of Environmental Protection
mg	milligram
µg	microgram
NAWQC	National Ambient Water Quality Criteria
ND	non-detect
NFG	National Functional Guideline
NOAA	National Oceanic and Atmospheric Administration
NOAEL	no observable adverse effect level
ns	not significant
<i>p</i>	probability
PAH	polycyclic aromatic hydrocarbon
PARCC	precision, accuracy, representativeness, completeness, comparability
PCE	tetrachloroethene
PEC	probable effect concentration
PERA	probabilistic ecological risk assessment
ORNL	Oak Ridge National Laboratory
QA/QC	quality assurance/quality control
QAPP	quality assurance program plan
Q-Q plot	quantile-quantile plot
QSAR	quantitative structure-activity relationship
RAGS	risk assessment guidance for Superfund
RBSL	risk-based screening level
RPD	relative percent difference
SAP	sampling and analysis plan
SLERA	screening-level ecological risk assessment
SLVHD	Salt Lake Valley Health Department
SQG	sediment quality guideline
SQuiRT	screening quick reference table
SVOC	semivolatile organic compound
TEC	threshold effect concentration
TIC	tentatively identified compound

TPH	total petroleum hydrocarbon
TPHCWG	Total Petroleum Hydrocarbon Criteria Working Group
TRV	toxicity reference value
UCL	upper confidence limit
UDEQ	Utah Department of Environmental Quality
UPL	upper prediction limit
USEPA	United States Environmental Protection Agency
VOC	volatile organic compound
WEF	wildlife exposure factor
WQS	water quality standard
WRS	Wilcoxon rank sum test

## EXECUTIVE SUMMARY

### E1.0 INTRODUCTION

On 12 June 2010, approximately 800 barrels (34,000 gallons) of crude oil was released from a Chevron pipeline (hereafter referred to as the Spill) into Red Butte Creek (in Salt Lake City, Utah) just downstream of the Red Butte Garden Arboretum (Chevron Pipeline Company [CPL] 2011). As of 9 September 2010, a total of 778 of the 800 barrels were accounted for through recovery from water, soil removal, and evaporation (CPL 2011). The purpose of this ecological risk assessment (ERA) is to evaluate the potential for adverse ecological impacts that may occur as a result of potential exposures to residual concentrations of spill-related petroleum hydrocarbons following remediation efforts in Lower Red Butte Creek. This assessment does not address potential ecological effects from the spill or remediation prior to August 2011. Methods used to conduct the ERA are consistent with the State of Utah and United States Environmental Protection Agency (USEPA) guidance. Findings of this ERA are intended to support evaluations/determinations of whether:

- The remediation response was sufficient to protect biota of concern;
- A more detailed ERA is warranted for this urban creek; and/or
- There is a need for additional risk management actions, and, if needed, what is the scope of these actions.

Consistent with guidance, a screening-level ERA (SLERA) was conducted and the elements of this SLERA included:

- Problem formulation;
- Exposure assessment;
- Effects assessment; and
- Risk characterization.

Key features of this SLERA include:

- Where applicable and possible, the SLERA is consistent in approach and methodology with the human health risk assessment (HHRA) that was performed concurrently.
- Utah Department of Environmental Quality (UDEQ 2005) *TPH Fractionation* guidance was used to evaluate total petroleum hydrocarbons (TPH). UDEQ's fractionation guidance builds on approaches previously described by the Total Petroleum Hydrocarbon Criteria Working Group (TPHCWG) and the Massachusetts Department of Environmental Protection (MaDEP). Where UDEQ

TPH benchmarks were lacking, MaDEP benchmarks were used to evaluate potential risks due to exposures to petroleum hydrocarbons—specifically, to aliphatic and aromatic carbon-chain fractions.

- Nearby urban creeks were identified to characterize local background conditions of urban creeks not impacted by the Spill and were established to correctly identify concentrations and biological responses attributable to the Spill. Local background creeks identified in the *Red Butte Creek Crude Oil Spill Water, Sediment and Macroinvertebrate Sampling Plan*, v. 17 (CPL 2011) and used in this ERA are Emigration Creek, Parleys Creek, City Creek, and Mill Creek.
- In-creek benthic community structure was evaluated as an added line of evidence to characterize the ecological significance of any identified ecological risks.

## E2.0

### DATA EVALUATION

In August 2011, bank soil, creek bed sediment, water, and benthic macroinvertebrate samples were collected in Lower Red Butte Creek and identified local background (urban) creeks in Salt Lake City, Utah. Data collection activities were designed to characterize (i) concentrations of petroleum-related constituents and (ii) structure of benthic macroinvertebrate communities for use in the risk assessment. For consistency and where possible to provide a baseline data set, all methods used to conduct the field collection and laboratory analyses were the same as those described in the *Red Butte Creek Crude Oil Spill Water, Sediment and Macroinvertebrate Sampling Plan* v. 17 [Incident Monitoring Sampling and Analysis Plan (SAP)] (CPL 2011).

Surface water, sediment, and bank soil samples were analyzed for:

- TPH;
- Volatile organic compounds (VOCs), including benzene, toluene, ethylbenzene, xylenes (BTEX); and
- Semivolatile organic compounds (SVOCs), including polycyclic aromatic hydrocarbons (PAHs); and
- Grain size and total organic carbon [for soil and sediment only].

Benthic macroinvertebrate samples were collected by Division of Water Quality personnel and analyzed by the National Aquatic Monitoring Center (also known as the “BugLab”) at Utah State University. To the extent possible, macroinvertebrate sampling locations were co-located

with surface water and sediment sampling stations to facilitate correlation of chemistry and biological data. Collection of macroinvertebrate community structure data is intended to provide additional evidence for characterizing/verifying potential ecological risks due to exposures to residual Spill-related petroleum hydrocarbons.

Data validation was conducted according to USEPA National Functional Guidelines (NFGs) (USEPA 1999a, 2004). Newer NFGs are available, but the SW-846 methods are better represented by the earlier versions of NFGs. All of the chemistry data were subject to a Level II review. A Level IV data validation was conducted on 10 percent of the data. Appropriate validation qualifiers were assigned to the data. Benthic macroinvertebrate data underwent the BugLab quality control procedures.

An organic compound was presumed not to exist in a particular environmental medium if it was never detected (100 percent non-detect) and detection limits met data quality objectives. Consistent with guidance (USEPA 1989), constituents that were detected at a frequency less than 5 percent were not quantitatively evaluated in this SLERA. The omission of a quantitative evaluation for these rarely detected constituents is discussed further in the uncertainty analysis.

### **E3.0 LOCAL BACKGROUND EVALUATION**

Nearby urban creeks were selected to represent in-creek local background conditions having similar environmental expectations as Lower Red Butte Creek in the absence of the effects of the Spill. As such, local background conditions can be used to characterize the "reasonable attainable" state and can provide the point-of-local background to assess the potential impairment. A Local Background Evaluation was performed by McDaniel-Lambert to support both the HHRA and ERA (McDaniel-Lambert 2012). The purpose of this evaluation was to determine whether concentrations of PAHs detected in samples within Lower Red Butte Creek are comparable to concentrations of PAHs detected in local background creeks.

There were insufficient data to conduct comparisons to local background creeks for:

- Bis(2-ethylhexyl)phthalate and bromoform in surface water. Bis(2-ethylhexyl)phthalate and bromoform are not considered petroleum-related constituents and are not evaluated further in this SLERA (consistent with the HHRA).

- 1-Methylnaphthalene, 2-methylnaphthalene, anthracene, benzo(a)anthracene, benzo(b)fluoranthene, benzo(g,h,i)perylene, benzo(k)fluoranthene, dibenzo(a,h)anthracene, fluorine, indeno(1,2,3-cd)pyrene, and naphthalene in creek bed sediments
- 1-Methylnaphthalene, 2-methylnaphthalene, dibenzo(a,h)anthracene, fluorene, and naphthalene in soil/sediment.

However, where there were at least six detections, findings of both two-sample and quantile tests suggest that concentrations of PAHs and TPH in Red Butte Creek were comparable or less than in local background creeks.

Visual examination of quantile-quantile (Q-Q) plots indicated inconsistencies with statistical tests for some PAHs and TPH-motor oil. However, the HHRA concluded that the number of samples was adequate to detect desired differences between Red Butte and the local background creeks (McDaniel-Lambert 2012). Further, individual site observations were compared against a background threshold value (BTV) to determine whether or not point-by-point site concentrations are within local background creek concentrations. The BTV analysis supports the conclusion that elevated site PAH concentrations occur in localized areas of Lower Red Butte Creek – namely 1731 East 900 South, Above 1500 East, and 1225 East Harvard Avenue. All other sampling locations are within expected local background creek levels.

The PAH composition analysis did not reveal any differences between Lower Red Butte Creek and background PAH ratios, including the PAH ratios for the maximum detections in Lower Red Butte Creek sediment. These findings suggest that petroleum-related hydrocarbons detected in Red Butte Creek appear to be consistent with PAHs typical of urban runoff.

The weight of evidence shows that most Lower Red Butte Creek PAHs are consistent with local background creek levels; however, a few creek locations appear to exceed background levels. However, for the purposes of this SLERA, comparisons to local background creek concentrations were not used to identify constituents of potential ecological concern and findings of this local background creek evaluation were discussed in the uncertainty analysis and conclusions.

## **E4.0**


### **PROBLEM FORMULATION**

Problem formulation establishes the scope of the ecological risk assessment, identifies the major factors to be considered, and ensures that



ecological receptors likely to be exposed and exposure scenarios most likely to contribute to ecological risk are evaluated. A conceptual site model (CSM) was prepared that identifies and summarizes the sources, mechanisms of transport, media of concern, exposure routes, and receptor groups and is intended to identify those exposure scenarios that are most likely to put biotic receptors of concern at risk. A CSM for the Lower Red Butte Creek ERA is shown in Figure E4-1.

Figure E4-1. Conceptual Site Model for the Lower Red Butte Creek SLERA



Note that no federal- or state-listed threatened and endangered species reside and no designated critical habitat was identified in the reach of interest for Lower Red Butte Creek.

Constituents of potential ecological concern (COPECs) are petroleum-related constituents that may adversely affect biota. Petroleum-related constituents include TPH, TPH fractions, and hazardous components of TPH (in particular, PAHs and BTEX) (Agency for Toxic Substances and Disease Registry 1999; TPHCWG 1997a). COPECs do not necessarily signify a risk; rather, they are merely constituents that have been identified for further examination. COPECs for each medium of concern are listed in Table E4-1.

Benzene and ethylbenzene were not detected in creek soils/sediments. Xylene was not detected in creek sediments, but was detected in creek soils/sediment; however, xylene was not detected at a concentration in creek soils/sediments greater than the ecological screening level (ESL) protective of wildlife. Although detected in 1 of 16 sediment samples, toluene was not detected at a concentration greater than ESLs for creek sediments. Note that acetone, chloroform, methylene chloride, tetrachloroethene (PCE), and bis(2-ethylhexyl)phthalate are not considered petroleum-related constituents and are not evaluated further in this SLERA.

**Table E4-1. Constituents of Potential Ecological Concern**

Constituent	Surface Water*	Creek Bed Sediment	Creek Soil/Sediment
<b>Low Molecular Weight PAHs</b>			
Anthracene		X	
<b>High Molecular Weight PAHs</b>			
Benzo(a)anthracene		X	
Benzo(a)pyrene		X	
Benzo(b)fluoranthene		X	
Benzo(g,h,i)perylene		X	
Benzo(k)fluoranthene		X	
Dibenzo(a,h)anthracene		X	
Indeno(1,2,3-cd)pyrene		X	
Pyrene		X	

Constituent	Surface Water*	Creek Bed Sediment	Creek Soil/Sediment
<b>TPH</b>			
TPH Diesel			
Aromatics		X	X
Aliphatics		X	X
TPH Motor Oil			
Aromatics		X	X
Aliphatics		X	X
<i>Notes:</i> X = COPEC * Benzene and ethylbenzene were not detected in creek soils/sediments. Although detected in 1 of 16 sediment samples, toluene was not detected at a concentration greater than the ESL for creek sediments. Although detected, acetone, bis(2-ethylhexyl)phthalate, bromoform, and PCE are not considered petroleum-related constituents and are not evaluated further in this SLERA (as consistent with the HHRA)			

## E5.0

## SUMMARY OF SCREENING-LEVEL ECOLOGICAL RISK ASSESSMENT FINDINGS

A key feature of this SLERA is the use of multiple lines of evidence (where available) to support characterizations of risk. The use of multiple lines of evidence (e.g., local background creek evaluation, risk estimates for local background creeks, benthic macroinvertebrate community metrics) is intended to provide several perspectives to assist in characterizing the potential for ecological risk.

**Aquatic Biota.** Potential risks to aquatic biota due to residual exposures of petroleum-related constituents in surface water appear to be negligible. All analytes, except for bis(2-ethylhexyl)phthalate, were either not detected or had maximum concentrations less than risk-based ESLs. Note that bis(2-ethylhexyl)phthalate was only detected in 1 of 16 surface water samples and this compound is not considered a petroleum-related constituent .

**Benthic Macroinvertebrate Community.** Potential risks to benthic macroinvertebrate communities due to residual exposures of petroleum-related constituents in creek bed sediments also appear to be limited. COPECs were limited to eight PAHs, TPH-diesel, and TPH-motor oil. Exposures for 6 of 8 PAHs were less than threshold effect concentrations (TECs), leading to a conclusion that adverse effects are not expected to occur (MacDonald et al. 2000). Exposures to anthracene were greater than TECs, but less than probable effect concentrations, and a determination of

toxicity or nontoxicity cannot be confidently predicted (MacDonald et al. 2000).

An evaluation of local background creeks found that concentrations of PAHs in creek bed sediments of Lower Red Butte Creek were comparable or less than concentrations in creek bed sediments of local background creeks. In addition, exposures used in this SLERA appear to be comparable between Lower Red Butte Creek and local background creeks. Finally, metrics suggest that the structure of the benthic macroinvertebrate community in Lower Red Butte Creek is comparable to local background creeks. These lines of evidence suggest that potential exposures/risks in Lower Red Butte Creek are comparable to conditions observed in local background creeks and are unlikely to be attributable to residual Spill-related petroleum hydrocarbons.

**Riparian Birds and Mammals.** Potential risks to riparian birds and mammals due to residual exposures to aromatic and aliphatic fractions of TPH in Lower Red Butte Creek were identified. Note that exposures and risk estimates for aromatic and aliphatic fractions of TPH-diesel were comparable to or less than those measured/calculated for the local background creeks. Although statistical tests found no significant difference, visual examination of Q-Q plots suggest that TPH-motor oil concentrations appear to be greater in Red Butte Creek as compared to local background creeks. Risk estimates for the aliphatic fraction of TPH-motor oil were greater than one and greater than risk estimates calculated for the local background creeks.

**Local Background Evaluation.** An evaluation of local background creeks found that concentrations of several PAHs and TPH-diesel in creek bed sediment of Lower Red Butte Creek were comparable or less than concentrations in creek bed sediment of local background creeks. Similarly, this evaluation found that concentrations of several PAHs and TPH-diesel in soil/sediment of Lower Red Butte Creek were comparable or less than concentrations in soil/sediments of local background creeks. Although visual examination of Q-Q plots found inconsistencies with the statistical analyses, an analysis of PAH composition suggests a source of PAHs that is consistent with urban runoff (McDaniel-Lambert 2012). This evaluation of the local background creek conditions suggest that petroleum hydrocarbons detected in Red Butte Creek may not be Spill-related.

**Uncertainties.** Uncertainties associated with the risk analyses were identified for this SLERA. To reduce uncertainties, focused verification of

this SLERA may be considered. However, given potential risks to biota in Lower Red Butte Creek, for the most part, appear to be comparable to or less than risks for conditions in local background creeks, the need for verification may not be considered essential to support decision-making.

• This page intentionally left blank •

## 1.0 INTRODUCTION

Red Butte Creek is a perennial third-order stream found in Salt Lake City, Utah. Red Butte Creek has a lengthy wild land reach in the Wasatch Front Range, and then flows through residential/urban reaches before entering the Jordan River. On 12 June 2010, approximately 800 barrels (34,000 gallons) of crude oil was released from a Chevron pipeline (hereafter referred to as the Spill) into Red Butte Creek just downstream of the Red Butte Garden Arboretum (Chevron Pipeline Company [CPL] 2011). Immediately following the Spill (summer/fall 2010), Phase 1 of the creek cleanup was initiated. Approximately 400 barrels were recovered at the Spill site on land and about 400 barrels entered Lower Red Butte Creek. As of 9 September 2010, a total of 778 of the 800 barrels were accounted for through recovery from water, soil removal, and evaporation (CPL 2011).

On behalf of the Utah Department of Environmental Quality (UDEQ) Division of Water Quality, the oversight agency of the Spill, ERM-West, Inc. (ERM) has prepared this Ecological Risk Assessment (ERA) to assess the potential risks to biota associated with residual oil<sup>1</sup> in Lower Red Butte Creek in Salt Lake City County, Utah.

## 1.1 SPILL HISTORY

Under the oversight of the Unified Command, CPL initiated cleanup, recovery, and restoration activities. Immediate measures were taken to minimize the impact of the crude oil. Remediation activities were carried out in accordance with the approved *Removal Action Plan* (ENACT 2010) developed by CPL, Salt Lake City, Salt Lake Valley Health Department (SLVHD), and UDEQ. Response measures included maintaining boom operations and deploying emergency response equipment. Phase 1 of the creek cleanup consisted of crews working the length of the creek (i) manually washing the creek to remove oil from rocks, sediments, and vegetation; and (ii) flushing the creek with high water flow. Efforts to restore conditions within the riparian corridor of this urban creek

---

<sup>1</sup> This assessment does not address potential ecological effects from the spill or remediation prior to August 2011.

included restoring vegetation, stabilizing banks, and re-introducing native trout.

## 1.2 OVERVIEW OF THE ECOLOGICAL RISK ASSESSMENT

The ERA process is used to systematically evaluate and organize data, assumptions, and uncertainties to help understand and predict the relationships between stressors and ecological effects in a way that is useful for environmental decision-making (United States Environmental Protection Agency [USEPA] 1989). The purpose of this ERA is to evaluate the potential for adverse ecological impacts that may occur as a result of potential exposures to residual concentrations of spill-related petroleum hydrocarbons following remediation efforts in Lower Red Butte Creek. Methods used to conduct the ERA are discussed in further detail in Appendix A of this report and are consistent with the following State of Utah and USEPA guidance:

- Utah Administrative Code, Rule R315-101-5, *Health Evaluation Criteria, Risk Assessment*;<sup>2</sup>
- *Framework for Ecological Risk Assessment* (USEPA 1992a);
- *Ecological Risk Assessment Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessments* (USEPA 1997); and
- *Guidelines for Ecological Risk Assessment* (USEPA 1998).

Findings of this ERA<sup>3</sup> are intended to support evaluations/determinations of whether:

- The remediation response was sufficient to protect creek biota;
- A more detailed ERA is warranted for this urban creek; and/or
- There is a need for additional risk management actions, and, if needed, what is the scope of these actions.

---

<sup>2</sup> Utah Administrative Code, Rule R315-101-5, *Health Evaluation Criteria, Risk Assessment* specifically applies to Hazardous Waste sites. Nonetheless, where possible, this ERA will be consistent with this State rule.

<sup>3</sup> The findings of the Local Background Evaluation (Section 4) were used to determine whether the source of the hydrocarbons detected in Lower Red Butte Creek were related to the Spill.



### 1.2.1 *Tiered Approach*

A tiered ERA approach is being employed as needed (Figure 1-1) and includes:

- Tier 1: Screening-Level ERA (SLERA);
- Tier 2: Baseline ERA (BERA); and
- Tier 3: Probabilistic ERA (PERA).

A scientific/management decision point exists at the conclusion of each tier, when it will be decided:

1. Whether or not the risk assessment, in its current state, is sufficient to support decision-making; and
2. If the assessment is determined to be insufficient, whether or not refinement of the current tier or progression to the next tier would provide a sufficient benefit to warrant the additional effort.

Advantages of a tiered approach include:

- Provides opportunities for regular input and direction by decision-makers;
- Provides a logical, stepwise approach for compiling and analyzing more site-specific information and incorporating more realistic assessments of exposure and effects; and
- Provides opportunities to streamline and focus the ERA-related effort.

### 1.2.2 *Elements and Key Features of this Ecological Risk Assessment*

This report focuses on the approach and findings of the SLERA conducted for Lower Red Butte Creek. Elements of this SLERA include (Figure 1-2):

- Problem formulation;
- Exposure assessment;
- Effects assessment; and
- Risk characterization.

Key features of this SLERA include:

- Where applicable and possible, the SLERA is consistent in approach and methodology with the human health risk assessment (HHRA) that is being performed concurrently.

- UDEQ (2005) *TPH Fractionation* guidance was used to evaluate total petroleum hydrocarbons (TPH). UDEQ's fractionation guidance builds on approaches previously described by the Total Petroleum Hydrocarbon Criteria Working Group (TPHCWG) and the Massachusetts Department of Environmental Protection (MaDEP). Where UDEQ TPH benchmarks were lacking, MaDEP benchmarks<sup>4</sup> were used to evaluate potential risks due to exposures to petroleum hydrocarbons – specifically, to aliphatic and aromatic carbon-chain fractions.
- Nearby urban background creeks were identified to characterize local background conditions of urban creeks not impacted by the Spill and were established to correctly identify concentrations and biological responses attributable to the Spill. Local background creeks identified in the *Red Butte Creek Crude Oil Spill Water, Sediment and Macroinvertebrate Sampling Plan*, v. 17 (CPL 2011) and used in this ERA are Emigration Creek, Parleys Creek, City Creek, and Mill Creek.<sup>5</sup>
- In-creek benthic community structure was evaluated as an added line of evidence to characterize the ecological significance of any identified ecological risks.

---

<sup>4</sup> MaDEP (2002, 2007)

<sup>5</sup> Emigration Creek, Parleys Creek, City Creek, and Mill Creek were not affected by the Spill and are considered to be representative of urban creeks in the Salt Lake City area (CPL 2011). Like Red Butte Creek, these local background creeks have lengthy wild land reaches in the Wasatch front range, and then flow through residential/urban reaches before entering the Jordan River.

## 2.0 *SITE DESCRIPTION*

Descriptions of Lower Red Butte Creek were primarily obtained from:

- *Salt Lake City Riparian Corridor Study: Final Red Butte Creek Management Plan* (Bio-West 2010);
- *Red Butte Creek Crude Oil Spill Water, Sediment and Macroinvertebrate Sampling Plan* (CPL 2011); and
- Utah Division of Wildlife Resources documents.<sup>6</sup>

This site description is intended to give a general sense of the setting in Lower Red Butte Creek. It is not intended to be a treatise on the abiotic/biotic features of Lower Red Butte Creek.

## 2.1 *LOCATION/GENERAL SETTING*

Red Butte Creek is a narrow rocky creek located between City Creek to the north and Emigration Creek to the south (CPL 2011; Bio-West 2010) (Figure 2-1). Red Butte Creek has been divided into Upper Red Butte Creek (upstream of Red Butte Gardens) and Lower Red Butte Creek (downstream of Red Butte Gardens) (Bio-West 2010). Upper Red Butte Creek drains approximately 5,400 acres of mountainous land primarily owned and managed by the U.S. Forest Service. Red Butte Reservoir and the Red Butte Creek Research Natural Area are located in Upper Red Butte Creek.

The reach impacted by the Spill, Lower Red Butte Creek, passes through an urban area where multiple point and nonpoint sources likely input to the creek. The open channel portion of Lower Red Butte Creek terminates at approximately 900 East where the creek enters a series of culverts that discharge to Liberty Park Lake at Liberty Park. The 1300 South conduit then conveys the flows from Lower Red Butte Creek and Emigration Creek to the Jordan River via a 3.4-mile-long pipe. One function of the impacted portion of Lower Red Butte Creek is as an urban storm water conveyance system. There are campus parking lots and roadways

---

<sup>6</sup> <http://wildlife.utah.gov/dwr/>

immediately adjacent to the spill site. The impacted reach drops about 750 feet over a reach of 18,000 feet, averaging approximately a 4 percent drop (Figure 2-2).

## 2.2 GEOLOGY

The surface geology of the Upper Red Butte Creek is composed of various members of the Triassic Ankareh formation as well as Jurassic/Triassic Nugget Sandstone (Bio-West 2010). Approximately 50 to 86 percent of the soils in the upper subwatershed have severe erosion potential. Lower Red Butte Creek flows through deposits ranging in size from finer-grained silt and clay to coarser sand and gravel deposits where 20 to 35 percent of the soil has severe to very severe erosion potential. Median streambed particle sizes range from 12 to 75 millimeters. Medium and large-sized gravel are the dominant substrate sizes in riffle areas of Lower Red Butte Creek.

## 2.3 IN-CREEK FLOWS

Red Butte Creek has a perennial flow upstream of Red Butte Reservoir and is considered to have “perennial-reduced” flow below that point (Bio-West 2010). Although flow is regulated by the Red Butte Reservoir, the creek’s hydrology is characterized by a distinct springtime peak in flow, which is typical of snowmelt systems. Flows in Lower Red Butte Creek are “flashy” with rapid, brief rises in flow during storms, a typical pattern followed by urban creeks. Average annual high flows are 22 cubic feet per second (cfs), while typical base flows are 2 cfs. Episodic high flows are likely to affect the transport/spatial distribution of chemicals, as well as physically affect biotic communities.

## 2.4 FLORA AND FAUNA

The most common trees along the streamside areas of Red Butte Creek are box elder (*Acer negundo*) and cottonwood (*Populus* sp.), with Gambel oak (*Quercus gambelii*) dominant in undeveloped upper slope areas. Siberian elm (*Ulmus pumila*), an introduced invasive tree species, is also fairly common. Russian olive (*Elaeagnus angustifolia*), an introduced invasive tree, is present, but less prominent. Common shrub species include redosier dogwood (*Cornus sericea*), twinberry honeysuckle (*Lonicera involucrata*), and narrowleaf willow (*Salix exigua*), with Woods’ rose (*Rosa woodsii*) common on upper portions of slopes. The understory vegetation layer includes native species such as Western poison ivy (*Toxicodendron rydbergii*) and Virginia creeper (*Parthenocissus quinquefolia*), and field

horsetail (*Equisetum arvense*). Introduced species such as ornamental English ivy (*Hedra helix*), common periwinkle (*Vinca minor*), climbing nightshade (*Solanum dulcamara*), smooth brome (*Bromus inermis*), and lesser burdock (*Arctium minus*) are significant components of the understory cover in several reaches. In addition, the upper slope portions of some reaches contain the invasive species whitetop (*Cardaria draba*) and houndstongue (*Cynoglossum officinale*). Canopy (tree) cover is generally high, though is markedly reduced in the lower urban reaches (Bio-West 2010).

A review of the Utah Natural Heritage Program's Biodiversity Tracking and Conservation System, Utah Division of Wildlife Resources, and U.S. Fish and Wildlife Service found (i) no federal- or state-listed threatened and endangered species; and (ii) no designated critical habitat residing in the reach of interest for Lower Red Butte Creek. A refuge population of endangered June sucker (*Chasmistes liorus*) currently inhabits Red Butte Reservoir (Bio-West 2010). However, Red Butte Reservoir is located upstream of the spill site, and there are no known occurrences of the June sucker in Lower Red Butte Creek.

A managed population of native Bonneville cutthroat trout (*Oncorhynchus clarki utah*) exists in the creek above the Red Butte Reservoir (Bio-West 2010). Lower Red Butte Creek is not reported in agency publications as supporting a fishery (Salt Lake County [SLCO] 2009), but trout have been observed in the creek, perhaps from private landowners stocking small numbers of trout for fishing (Bio-West 2010). The Utah Division of Wildlife Resources stocked 3,000 fingerling cutthroat trout in November 2011.

Bio-West (2010) concludes that limited information is available about the wildlife of the urban lower portion of Red Butte Creek. Deer, raccoon, and skunk have been observed along the Lower Red Butte Creek. During the Audubon Society's 2005 Christmas bird count, over 30 different species of birds were observed within the University of Utah survey area, which includes portions of the Lower Red Butte Creek riparian corridor (Bio-West 2010). Miller Bird Refuge and Bonneville Glen Park are generally recommended for recreational bird watching.

• This page intentionally left blank •

### **3.0 DATA EVALUATION**

The following subsections describe data collection, data validation, and data usability in support of this ERA, as well as the handling of non-detected values.

#### **3.1 DATA COLLECTION**

In August 2011, bank soil, creek bed sediment, water, and benthic macroinvertebrate samples were collected in Lower Red Butte Creek and identified local background (urban) creeks in Salt Lake City, Utah. Data collection activities were designed to characterize (i) concentrations of petroleum-related constituents and (ii) structure of benthic macroinvertebrate communities for use in the risk assessment.

For consistency and where possible to provide a baseline data set, all methods used to conduct the field collection and laboratory analyses were the same as those described in the *Red Butte Creek Crude Oil Spill Water, Sediment and Macroinvertebrate Sampling Plan* v. 17 [Incident Monitoring Sampling and Analysis Plan (SAP)] (CPL 2011).

##### **3.1.1 Sampling Locations**

Sampling stations were added to supplement sampling stations identified in the Incident Monitoring SAP (CPL 2011) to provide sufficient sample numbers in support of the SLERA (Table 3-1).

**Table 3-1. Incident Monitoring Sampling and Analysis Plan and Supplemental Sampling Stations to Support the Ecological Risk Assessment**

Site	Surface Water & Sediment Chemistry				Macroinvertebrate			
	SAP <sup>a</sup>	ERA <sup>b</sup>	# Samples		SAP <sup>a</sup>	ERA <sup>b</sup>	# Samples	
			Upstr <sup>c</sup>	Urban <sup>d</sup>			Upstr <sup>c</sup>	Urban <sup>d</sup>
Lower Red Butte Creek	7	6 <sup>e</sup>	1	12	3	6	1	8
Local Background Creeks								
Emigration Creek	1	3	1	3	2	1	1	2
Parley's Creek	1	3	0	4	0	2	0	2
Mill Creek	1	3	1	3	2	1	1	2
City Creek	1	3	2	2	1	2	1	2
Total Lower Red Butte Creek <sup>d</sup>				12				8
Total Local Background Creeks <sup>d</sup>				12				8

*Notes:*

- a. Sampling locations from the Incident Monitoring SAP (CPL 2011).
- b. Sampling locations added to support the ERA.
- c. Sampling locations in upstream (of spill) or natural reach of creeks.
- d. Sampling locations in urbanized reach of creeks.
- e. Assumes that the same sample can be used to support both the HHRA and ERA.

### 3.1.1.1 Lower Red Butte Creek Sampling Locations

Bank soil, creek bed sediment, surface water, and benthic macroinvertebrate samples were collected within Lower Red Butte Creek (Table 3-1). The 12 downstream locations were selected to characterize the potential nature and extent of residual Spill-related impacts within Lower Red Butte Creek (CPL 2011) (Figure 3-1; Table 3-1).

### 3.1.1.2 Local Background Creek Sampling Locations

Petroleum hydrocarbons are commonly detected in urban storm water. Hence, to establish relevant local background concentrations of petroleum hydrocarbons and to assess potential risks attributable to the Spill, bank soil, creek bed sediment, surface water, and benthic macroinvertebrate samples were also collected for the following local background creeks<sup>7</sup> selected for Lower Red Butte Creek (Figure 3-1; Table 3-1):

<sup>7</sup> Emigration Creek, Parleys Creek, City Creek, and Mill Creek are considered representative of the local background levels of hydrocarbons present in Lower Red Butte Creek (CPL 2011).



- Emigration Creek
- Parleys Creek
- City Creek
- Mill Creek

Like Red Butte Creek, these local background creeks have lengthy wild land reaches in the Wasatch Front Range, and then flow through residential/urban reaches before entering the Jordan River (Figure 2-1).<sup>8</sup> Emigration Creek, Parleys Creek, City Creek, and Mill Creek were not affected by the Spill and are considered to be representative of local background levels of hydrocarbons present in urban creeks in the Salt Lake City area (CPL 2011).

Photographs of Lower Red Butte Creek and local background creeks are provided in Appendix B.

### 3.1.2 *Media and Analytes of Interest*

Surface water, sediment, and bank soil samples were analyzed for:

- TPH;
- Volatile organic compounds (VOCs), including benzene, toluene, ethylbenzene, xylenes (BTEX); and
- Semivolatile organic compounds (SVOCs), including polycyclic aromatic hydrocarbons (PAHs); and
- Grain size and total organic carbon [for soil and sediment only].

### 3.1.3 *Benthic Macroinvertebrates*

Benthic macroinvertebrate samples were collected by Division of Water Quality personnel and analyzed by the National Aquatic Monitoring Center (also known as the “BugLab”) at Utah State University. To the extent possible, macroinvertebrate sampling locations were co-located with surface water and sediment sampling stations to facilitate correlation of chemistry and biological data. Collection of macroinvertebrate community structure data is intended to provide additional evidence for

---

<sup>8</sup> The identified local background creeks have land uses and land covers roughly similar to Red Butte Creek.

characterizing/verifying potential ecological risks due to exposures to residual Spill-related petroleum hydrocarbons. The following community metrics were computed and provided by the BugLab at Utah State University:

- Taxa richness
- Evenness
- Species diversity
- Hilsenhoff biotic index (HBI)
- Percent chironomids<sup>9</sup>
- Percent EPT<sup>10</sup>

For further discussion regarding the use and interpretation of these community metrics, please see Sections 7.4 and 8.3.

## 3.2 DATA VALIDATION

Data validation was conducted according to USEPA National Functional Guidelines (NFGs) (USEPA 1999a, 2004). Newer NFGs are available, but they are guidelines for USEPA's Contract Laboratory Program methods. The SW-846 methods are better represented by the earlier versions of NFGs.

### 3.2.1 Analytical Chemistry Data Quality Control

All of the chemistry data were subject to a Level II review. A Level II review consists of a review of all sample-related quality control parameters, including holding times, blank contamination, laboratory control sample, matrix spike/matrix spike duplicate, and surrogates.

In addition, a Level IV data validation was conducted on 10 percent of the data. Level IV data validation consisted of a review of all parameters reviewed as part of the Level II review with additional review of instrument performance check (as applicable), initial and continuing calibrations, and internal standards (as applicable). In addition, Level IV includes review of the raw data, including chromatograms, log books, quantitation reports, and spectra.

---

<sup>9</sup> midge—considered to be a pollutant-tolerant taxa (USEPA 1999b).

<sup>10</sup> ephemeroptera (mayfly), plecoptera (stonefly), and trichoptera (caddisfly)—considered to be pollutant-sensitive taxa (USEPA 1999b).

Appropriate validation qualifiers were assigned to the data. All of the data, including qualified data, were considered usable and no data were rejected. The Quality Assurance/Quality Control (QA/QC) memorandums are included as Appendix C.

### 3.2.2 *Benthic Macroinvertebrate Data Quality Control*

Benthic macroinvertebrate data underwent the BugLab quality control procedures. The following three potential sources of error were carefully checked to assure quality of the data:

- Sample sorting;
- Macroinvertebrate identification; and
- Data processing.

**Sample Sorting.** Fifteen percent of all sorted samples were examined to ensure that at least 95 percent of the organisms were removed from the examined material. A sorting effectiveness ( $E_s$ ) number was computed as:

$$E_s = 100 * S / (R + S) \quad \dots \text{Eq 3-1}$$

Where R is the total number of organisms obtained during the resort of the remnant material and S is the total number of organisms originally obtained from the sample sorting. All samples exceeded the goal of greater than or equal to 95 percent of the organisms removed during the original sort. The average sorting effectiveness was 96.5 percent.

**Macroinvertebrate Identification.** To assure consistency in sample identification, at least 10 percent of the samples were re-identified by a second taxonomist. The identifications performed by the taxonomists were 99.2 percent similar as measured by the Bray-Curtis similarity index.

**Data Processing.** Ten percent of samples within this data set were checked against the bench sheets to ensure the accuracy of all of the information.

## 3.3 *DATA USABILITY*

The primary objective of the data review and usability evaluation was to identify appropriate data for use in the risk assessment. The analytical data were reviewed for applicability and usability following procedures in the *Guidance for Data Usability in Risk Assessment (Part A)* (USEPA 1992b, and USEPA 1989). According to USEPA Data Usability Guidance, there

are six principal evaluation criteria by which data are judged for usability in risk assessment. The six criteria are:

- Availability of information associated with site data;
- Documentation;
- Data sources;
- Analytical methods and detection limits;
- Data review; and
- Data quality indicators, including precision, accuracy, representativeness, comparability, and completeness.

A summary of the findings of the data usability effort in support of the ERA is provided in the Data Usability Worksheet from the *Risk Assessment Guidance for Superfund: Volume I Human Health Evaluation Manual – Part D* (USEPA 2001a). The Data Usability Worksheet summarizes the criteria used to identify data usability and is presented in Table 3-2.

**Table 3-2. Criteria Used to Identify Data Usability**

Activity	Comment
<b>Field Sampling</b>	
Discuss sampling problems and field conditions that affect data usability.	No field conditions resulted in poor sample recovery.
Are samples representative of receptor exposure for this medium (e.g. sample depth, grab vs. composite, filtered vs. unfiltered, low flow, etc.)?	Water samples were grab (discrete) samples. All sediment/bank soil samples were composite samples. All samples were representative of receptor exposures and analyzed for a broad spectrum of analyses.
Were samples appropriately documented and can they be correlated to a specific geographic location?	All samples reported by the laboratory were documented on the chain-of-custodies and were correlated to a specific geographic location.
Assess the effect of field quality control (QC) results on data usability.	No quality assurance program plan (QAPP) was available. Field, equipment and trip blanks, and soil duplicate samples were collected during all field sampling activities. No qualifications were made based upon field QC results.
Summarize the effect of field sampling issues on the risk assessment, if applicable.	There were no field sampling issues that affected the data quality for risk assessment purposes.
<b>Analytical Techniques</b>	
Were the analytical methods appropriate for quantitative risk assessment?	Yes, the analytical techniques used were appropriate for risk assessment purposes. Analytical techniques followed USEPA guidelines.

Activity	Comment
Were detection limits adequate?	Yes, detection limits were adequate for risk assessment purposes.
Summarize the effect of analytical technique issues on the risk assessment, if applicable.	There were no issues raised that were particular to the analytical techniques used. Analytical techniques for soils and water followed USEPA-based guidelines.
<b>Data Quality Indicators</b>	
Precision - How were duplicates handled?	The field duplicate samples were compared for consistency (relative percent difference [RPD] was calculated) and the maximum detection or the minimum reporting limit was selected. Several chemicals were analyzed by multiple analyses (e.g. PAHs by 8270SIM and 8270). In a few cases, the laboratory result for PAHs (USEPA 8270SIM) exceeded the calibration range. In that case, the laboratory indicated to use the result from the USEPA 8270C analysis.
Accuracy - How were duplicate samples handled?	No samples were qualified due to accuracy-related outliers such as an LCS, MS/MSD or surrogate recovery.
Representativeness - Indicate any problems associated with data representativeness (e.g., trip blank or rinsate blank contamination, chain of custody problems, etc.).	Chain of custody forms were checked by QC staff and laboratory was informed of any problems within 1 to 2 days of sample collection. Based on the procedures used, the data was representative of site conditions.
Completeness - Indicate any problems associated with data completeness (e.g., incorrect sample analysis, incomplete sample records, problems with field procedures, etc.).	There were no problems identified.
Comparability - Indicate any problems associated with data comparability.	USEPA methods were utilized throughout the project. Some analyses were conducted by Lancaster Laboratories and America West Laboratories. A subset of data was sent to Lancaster as QC samples. No issues were identified.
Were the data quality objectives (DQOs) specified in the QAPP satisfied?	No QAPP was available.
Summarize the effect of DQO issues on the risk assessment, if applicable.	Precision/Accuracy/Representativeness/Completeness/Comparability (PARCC) criteria met DQOs and resulted in usable data for the risk assessment.

Activity	Comment
<b>Data Validation and Interpretation</b>	
What are the data validation requirements?	For chemistry data, all laboratory reports were provided as either a Level II or Level III+ (Level IV equivalent) data package. The detailed data validation procedures are consistent with the USEPA National Functional Guidelines for Data Validation. The data were reviewed against the USEPA National Functional Guidelines.
What method or guidance was used to validate the data?	The USEPA National Functional Guidelines for Data Validation were used.
Was the data validation method consistent with guidance? Discuss any discrepancies.	Yes, data validation methods were consistent with the guidance.
Were all data qualifiers defined? Discuss those which were not.	Yes, all definitions of all data qualifiers are presented in the laboratory reports.
Which qualifiers represent usable data?	All data collected and validated are usable as qualified unless they are rejected with an R symbol.
Which qualifiers represent unusable data?	Data qualified as "R" (rejected) represents unusable data. No data were rejected.
How are tentatively identified compounds (TICs) handled?	TICs were reviewed and primarily consist of ketones, amides, amines, silanols, fatty acids, plant sterols and long chain alkanes. Amides, silanols, and amines are primarily linked to column breakdown. The long chain alkanes are potentially petroleum-related, however, some could be plant related hydrocarbons. A single detection of pyrene was identified as a TIC in the 8260 analysis of the sample named Red Butte Creek @ 1731 E. 900 S.–Bank. However, pyrene, is a target analyte of the 8270 SIM analysis and was detected in that same sample. Hence, TICs result in only a small source of uncertainty to the risk assessment.
Summarize the effect of data validation and interpretation issues on the risk assessment, if applicable.	Valid data were sufficient to perform the risk assessment. All data collected and validated are usable for the risk assessment as qualified.

### 3.4 HANDLING NON-DETECTED VALUES

Non-detects (NDs) or "left censored" data are inevitable in many environmental data sets. An organic compound was presumed not to exist in a particular environmental medium if it was never detected (100 percent ND) and detection limits met data quality objectives. Consistent with guidance (USEPA 1989), constituents that were detected

at a frequency less than 5 percent were not quantitatively evaluated in this ERA. The omission of a quantitative evaluation for these rarely detected constituents is discussed further in the uncertainty analysis (Section 8.6).

When greater than 5 percent of the data were comprised of non-detected concentrations, NDs were handled in accordance with guidance (USEPA 2006, 2010) and are described further in Appendix D. Appendix D also presents (i) descriptive statistics, including the 95 percent upper confidence limits (95UCLs); and (ii) comparisons between Lower Red Butte Creek and local background creeks.

• This page intentionally left blank •



#### 4.0 LOCAL BACKGROUND EVALUATION

Nearby urban creeks were selected to represent in-creek local background conditions having similar environmental expectations as Lower Red Butte Creek in the absence of the effects of the Spill. As such, local background creeks can be used to characterize the "reasonable attainable" state and can provide the point-of-reference to assess the potential impairment.

A Local Background Evaluation was performed by McDaniel-Lambert to support both the HHRA and ERA (McDaniel-Lambert 2012) (see Appendix D). In this Local Background Evaluation, PAH concentrations in Lower Red Butte Creek were compared to local background creeks for each of the following media of concern:

<u>Media of Concern:</u>	<u>Evaluate Exposures For:</u>
Surface water	Aquatic biota
Creek bed sediments	Benthic macroinvertebrates
Creek soil/sediment <sup>11</sup>	Riparian birds and mammals

The purpose of this evaluation was to determine whether concentrations of PAHs detected in samples within Lower Red Butte Creek are comparable to concentrations of PAHs detected in local background creeks.

A weight of evidence was used in this evaluation, including statistical tests, background threshold values (BTVs), and PAH diagnostic ratio comparisons. The exploratory analyses and comparative methods used are based on USEPA guidance (USEPA 2002). All statistical comparisons were conducted using USEPA's ProUCL v. 4.01.00. A more detailed discussion of this evaluation is found in the *Draft Human Health Risk Assessment* (McDaniel-Lambert 2012). Findings are briefly discussed below.

---

<sup>11</sup> While riparian wildlife may be exposed to both creek bank and creek bed substrate (combined bank soil and creek bed sediment), benthic macroinvertebrates were considered to be exposed only to creek bed (in-creek) sediments.

## 4.1 STATISTICAL COMPARISONS

Concentrations of petroleum-related hydrocarbons in Lower Red Butte Creek were compared to concentrations in local background creeks using one of the following two-sample statistical tests: (i) parametric *t*-test, or (ii) nonparametric Gehan test. In accordance with USEPA guidance (2002, 2010), the appropriate test was selected based on (McDaniel-Lambert 2012):

- Sample size (tests used only when sample size is eight or greater);
- Distribution of the data (normal or not normal);
- Equal or unequal variances; and
- Percent detected values.

These 2-sample tests evaluate the hypothesis:

H<sub>0</sub>: Mean/median concentrations in Lower Red Butte Creek are less than mean/median concentrations in local background creek.

To supplement the two-sample tests, the quantile test was conducted to evaluate the hypothesis (USEPA 2002):

H<sub>0</sub>: High concentrations in Lower Red Butte Creek are less than high concentrations in local background creeks.

**Surface Water.** Two constituents were detected in 1 of 17 surface water samples: bis(2-ethylhexyl)phthalate and bromoform. Bis(2-ethylhexyl)phthalate and bromoform are not considered petroleum-related constituents and will not be further evaluated in this SLERA (as consistent with the HHRA).

**Creek Bed Sediments.** The following nine PAHs that were detected in Lower Red Butte Creek were not evaluated because there were insufficient detected concentrations (i.e., less than four detected concentrations):

- |                        |                          |
|------------------------|--------------------------|
| • 1-Methylnaphthalene  | • Benzo(k)fluoranthene   |
| • 2-Methylnaphthalene  | • Dibenzo(a,h)anthracene |
| • Benzo(a)anthracene   | • Indeno(1,2,3-cd)pyrene |
| • Benzo(b)fluoranthene | • Naphthalene            |
| • Benzo(g,h,i)perylene |                          |

Findings of statistical tests for the remaining PAHs, TPH-diesel, and TPH-motor oil suggest (from Table 5 of McDaniel-Lambert 2012):

- There is no difference in median (central tendency) concentrations of PAHs detected in creek bed sediments between Lower Red Butte Creek and local background creeks.<sup>12</sup>
- There is no difference in high concentrations of PAHs detected in creek bed sediments between Lower Red Butte Creek and local background creeks.

**Creek Soil/Sediment.** Five PAHs that were detected in Lower Red Butte Creek were not evaluated because there were insufficient detected concentrations – i.e., less than four detected concentrations:

- 1-Methylnaphthalene
- 2-Methylnaphthalene
- Dibenzo(a,h)anthracene
- Fluorene
- Naphthalene

Findings of statistical tests for the remaining PAHs, TPH-diesel, and TPH-motor oil suggest (from Table 2 of McDaniel-Lambert 2012):

- There is no difference in median (central tendency) concentrations of PAHs detected in soil/sediment between Lower Red Butte Creek and local background creeks.<sup>13</sup>
- There is no difference in high concentrations of PAHs detected in soil/sediment between Lower Red Butte Creek and local background creeks.

## 4.2

### **COMPARISON TO BACKGROUND THRESHOLD VALUES**

Visual examination of quantile-quantile (Q-Q) plots indicated inconsistencies with statistical tests for some PAHs and TPH-motor oil. However, the HHRA concluded that the number of samples was adequate to detect desired differences between Red Butte and the reference creeks (McDaniel-Lambert 2012). Further, individual location observations in

---

<sup>12</sup> Do not reject the H<sub>0</sub> hypothesis.

<sup>13</sup> Do not reject the H<sub>0</sub> hypothesis.

Red Butte Creek for soil/sediment were compared against BTVs<sup>14</sup> calculated from local background creek data to determine whether specific location concentrations are within local background creek levels. Unlike the soil/sediment data, the creek bed sediment data did not show excessive evidence of potential hotspots with higher concentrations for the PAHs evaluated – therefore, a BTV analysis was determined to be unnecessary (McDaniel-Lambert 2012).

The BTV analysis for soil/sediment suggests that elevated site PAH concentrations may occur in localized areas of Lower Red Butte Creek – namely 1731 East 900 South, Above 1500 East, and 1225 East Harvard Avenue. All other sampling locations are within expected local background creek levels (Table 3 of McDaniel-Lambert 2012).

#### 4.3

#### ***POLYCYCLIC AROMATIC HYDROCARBONS COMPOSITION ANALYSIS***

A PAH composition analysis was conducted to determine which of the primary sources of PAHs commonly found in urban waterways might be the source of PAHs in Lower Red Butte Creek. These primary sources include:

- Pyrogenic: hydrocarbon compounds associated with the combustion of petroleum, wood, coal, etc.;
- Petrogenic: hydrocarbon compounds associated with petroleum, including fuel oils, coal, and lubricants; and
- Biogenic: associated with plant matter.

The following double-ratio cross plots were used as an exploratory tool to distinguish between pyrogenic and petrogenic sources:

- Benzo(a)anthracene to chrysene versus fluoranthene to pyrene; and
- Benzo(a)anthracene to benzo(a)pyrene versus fluoranthene to benzo(a)pyrene.

---

<sup>14</sup> BTVs were defined as the 95<sup>th</sup> percentile upper prediction limits (UPLs) (McDaniel-Lambert 2012).

This analysis shows overlap among the Lower Red Butte Creek and local background creek samples, with no distinct clustering. These findings reveal no differences in relative composition of PAHs and suggest that PAH sources in Lower Red Butte Creek and local background creeks are similar (i.e., urban runoff) (McDaniel-Lambert 2012).

#### 4.4

#### SUMMARY

There were insufficient data to conduct local background creek comparisons for:

- Bis(2-ethylhexyl)phthalate and bromoform in surface water;<sup>15</sup>
- 1-Methylnaphthalene, 2-methylnaphthalene, anthracene, benzo(a)anthracene, benzo(b)fluoranthene, benzo(g,h,i)perylene, benzo(k)fluoranthene, dibenzo(a,h)anthracene, fluorine, indeno(1,2,3-cd)pyrene, and naphthalene in creek bed sediments; and
- 1-Methylnaphthalene, 2-methylnaphthalene, dibenzo(a,h)anthracene, fluorene, and naphthalene in soil/sediment.

However, where there were at least six detections, findings of both two-sample and quantile tests suggest that concentrations of PAHs and TPH in Red Butte Creek were comparable or less than in local background creeks.

Visual examination of Q-Q plots indicated inconsistencies with statistical tests for some PAHs and TPH-motor oil. However, the HHRA concluded that the number of samples was adequate to detect desired differences between Red Butte and the local background creeks (McDaniel-Lambert 2012). Further, individual site observations were compared against a BTV to determine whether or not point-by-point site concentrations are within local background creek concentrations. The BTV analysis supports the conclusion that elevated site PAH concentrations occur in localized areas of Lower Red Butte Creek – namely 1731 East 900 South, Above 1500 East, and 1225 East Harvard Avenue. All other sampling locations are within expected local background creek levels.

---

<sup>15</sup> Bis(2-ethylhexyl)phthalate and bromoform are not considered petroleum-related constituents and are not evaluated further in this SLERA (consistent with the HHRA).

The PAH composition analysis did not reveal any differences between Lower Red Butte Creek and background PAH ratios, including the PAH ratios for the maximum detections in Lower Red Butte Creek sediment. These findings suggest that petroleum-related hydrocarbons detected in Red Butte Creek appear to be consistent with PAHs typical of urban runoff.

The weight of evidence shows that most Lower Red Butte Creek PAHs are consistent with local background creek levels; however, a few creek locations appear to exceed background levels. However, for the purposes of this SLERA, comparisons to local background creek concentrations were not used to identify constituents of potential ecological concern (COPECs) (Section 5.2) and findings of this Local Background Evaluation will be discussed in the uncertainty analysis and conclusions.

## 5.0 **PROBLEM FORMULATION**

Problem formulation establishes the scope of the ecological risk assessment, identifies the major factors to be considered, and ensures that ecological receptors likely to be exposed and exposure scenarios most likely to contribute to ecological risk are evaluated.

As described in the ERA work plan (Appendix A), problem formulation consists of the following:

- Identify biotic receptors of concern (BROC);
- Identify COPECs;
- Identify potentially complete exposure pathways; and
- Establish assessment endpoints.

## 5.1 **BIOTIC RECEPTORS OF CONCERN**

Given the number of species and the complexity of biological communities, all species present in Lower Red Butte Creek cannot be individually assessed. BROCs were identified to (1) focus the ERA on those receptors of concern, and (2) develop specific assessment endpoint statements.

Consistent with guidance (USEPA 1998), BROCs were identified and consider:

- Biota of regulatory interest – species and habitats that are protected by federal and state regulations;
- Biota of commercial/recreational interest – species that have an economic or recreational value (e.g. crops, livestock, fisheries, hunted game);
- Biota of resource management interest or habitats/species that may support functional attributes (e.g., flood control); and
- Biota of ecological interest – species that play an important role in mediating processes or interactions that affect the structure/function,

or biodiversity of native habitats, communities, or ecosystems (e.g., keystone species).<sup>16</sup>

All trophic levels, including primary producers, were considered.

***Biota of Regulatory Interest.*** No federal- or state-listed threatened and endangered species reside and no designated critical habitat was identified in the reach of interest for Lower Red Butte Creek. A refuge population of endangered June sucker (*Chasmistes liorus*) currently inhabits Red Butte Reservoir (Bio-West 2010). However, Red Butte Reservoir is located upstream of the Spill site, and there are no known occurrences of the June sucker in Lower Red Butte Creek.

***Biota of Commercial/Recreational Interest.*** Lower Red Butte Creek is not reported in agency publications as supporting a fishery (SLCO 2009), but trout have been observed in the creek, perhaps from private landowners stocking small numbers of trout for fishing (Bio-West 2010).

***Biota of Ecological Interest.*** Members of the following guilds were considered to play a key role in maintaining the structure/function of in-creek and riparian habitats and these guilds were identified as BROCs:

In-Creek Biota

- Aquatic plants
- Aquatic invertebrates
- Benthic macroinvertebrates
- Fish
- Amphibians<sup>17</sup>

Riparian Biota

- Reptiles<sup>13</sup>
- Waterfowl/shorebirds
- Mammals

---

<sup>16</sup> Plants and animals that provide shelter and/or food for special status species were also considered when identifying receptors of ecological concern.

<sup>17</sup> Given the lack of relevant widely accepted toxicity benchmarks, no quantitative evaluation of amphibians or reptiles will be conducted. The lack of a quantitative evaluation for amphibians and reptiles will be qualitatively discussed in the uncertainty analysis (Section 8.6).



The SLERA for Lower Red Butte Creek focuses largely on in-creek biota and riparian wildlife. As toxicity benchmarks used in this SLERA are inclusively protective of aquatic plants, aquatic invertebrates, and fishes, these receptors will be combined and evaluated as *Aquatic Biota*.

## 5.2 CONSTITUENTS OF POTENTIAL ECOLOGICAL CONCERN

For this SLERA, COPECs are petroleum-related constituents<sup>18</sup> that may adversely affect biota. COPECs do not necessarily signify a risk; rather, they are merely constituents that have been identified for further examination. COPECs were identified for the following media of concern:

<u>Media of Concern</u>	<u>Evaluate Exposures To</u>
• Surface water	• Aquatic biota
• Creek bed sediments	• Benthic macroinvertebrates
• Creek soil/sediment	• Riparian birds and mammals

A constituent was identified as a COPEC in Lower Red Butte Creek unless either of the following lines of evidence was true:

- Detected in less than 5 percent of the samples; or
- Maximum detected concentration is less than the corresponding risk-based ecological screening level (ESL).

Risk-based ESLs used in this SLERA are listed in Table 5-1 (see also Appendix A for further details).

**Table 5-1. Risk-Based Ecological Screening Levels**

(for constituents with greater than 5 percent frequency of detection)

<b>Constituent</b>	<b>Aquatic Biota</b> (µg/L)	<b>Sediment Biota</b> (mg/kg)	<b>Riparian Wildlife</b> (mg/kg)
<b>BTEX</b>			
Toluene	–	0.01	5.5
Xylene	–	–	10

<sup>18</sup> Petroleum-related constituents include TPH, TPH fractions, and hazardous components of TPH (in particular, PAHs and BTEX) (ATSDR 1999; TPHCWG 1997a).

Constituent	Aquatic Biota ( $\mu\text{g/L}$ )	Sediment Biota ( $\text{mg/kg}$ )	Riparian Wildlife ( $\text{mg/kg}$ )
<b>Low Molecular Weight PAHs</b>			
1-Methylnaphthalene	–	0.18	100
2-Methylnaphthalene	–	0.18	–
Anthracene	–	0.057	100
Fluorene	–	0.077	–
Naphthalene	–	0.18	100
Phenanthrene	–	0.20	100
<b>High Molecular Weight PAHs</b>			
Benzo(a)anthracene	–	0.11	1.1
Benzo(a)pyrene	–	0.15	1.1
Benzo(b)fluoranthene	–	0.027	1.1
Benzo(g,h,i)perylene	–	0.17	1.1
Benzo(k)fluoranthene	–	0.027	1.1
Chrysene	–	0.17	1.1
Dibenzo(a,h)anthracene	–	0.033	1.1
Fluoranthene	–	0.42	1.1
Indeno(1,2,3-cd)pyrene	–	0.017	1.1
Pyrene	–	0.20	1.1
<b>TPH</b>			
<b>TPH Diesel</b>			
Aromatics	–	0.090	na
Aliphatics	–	3.2	na
<b>TPH Motor Oil</b>			
Aromatics	–	na	na
Aliphatics	–	9.9	na
<i>Notes:</i> – = Less than 5 percent frequency of detection. $\mu\text{g/L}$ = Micrograms per liter $\text{mg/kg}$ = Milligrams per kilogram na = Not available			

COPECs lacking ESLs are identified and discussed as a part of the uncertainty analysis (Section 8.6).

Benzene and ethylbenzene were not detected in creek soils/sediments. Xylene was not detected in creek sediments, but was detected in creek soils/sediment; however, xylene was not detected at a concentration in creek soils/sediments greater than the ESL protective of wildlife. Although detected in 1 of 16 sediment samples, toluene was not detected at a concentration greater than the ESL for creek sediments.

Note that acetone, chloroform, methylene chloride, tetrachloroethene (PCE), and bis(2-ethylhexyl)phthalate were detected, are not considered petroleum-related constituents, and are not evaluated further in this SLERA. COPECs for each medium of concern are listed in Table 5-2. Further details related to the selection of COPECs are found in Appendix E.

**Table 5-2. Constituents of Potential Ecological Concern**

Constituent	Surface Water*	Creek Bed Sediment	Creek Soil/Sediment
<b>Low Molecular Weight PAHs</b>			
Anthracene		X	
<b>High Molecular Weight PAHs</b>			
Benzo(a)anthracene		X	
Benzo(a)pyrene		X	
Benzo(b)fluoranthene		X	
Benzo(g,h,i)perylene		X	
Benzo(k)fluoranthene		X	
Dibenzo(a,h)anthracene		X	
Indeno(1,2,3-cd)pyrene		X	
Pyrene		X	
<b>TPH</b>			
<b>TPH Diesel</b>			
Aromatics		X	X
Aliphatics		X	X
<b>TPH Motor Oil</b>			
Aromatics		X	X
Aliphatics		X	X
<i>Notes:</i> X = COPEC * For sediment-dwelling biota only. Acetone, bis(2-ethylhexyl)phthalate, bromoform, and PCE are not considered petroleum-related constituents, and are not evaluated further in this SLERA (consistent with the HHRA).			

### 5.3 POTENTIALLY COMPLETE EXPOSURE PATHWAYS

Potentially complete exposure pathways consist of:

- A source and mechanism of constituent release;
- A transport medium (e.g., soil, water, tissue);

- A point or area where receptors of concern may contact petroleum hydrocarbons (media concern); and
- An exposure route through which petroleum hydrocarbon uptake occurs (e.g., ingestion, inhalation, or dermal contact, including immersion).

Exposure routes that were considered include:

#### *In-Creek Biota*

- Direct contact (uptake) by aquatic biota for constituents in surface water; and
- Direct contact (uptake) by benthic macroinvertebrate biota for constituents in sediment.

#### *Riparian Biota*

- Direct (dermal) contact by wildlife for constituents in surface water and creek soil/sediment;
- Inhalation by wildlife for volatile constituents in surface water and creek soil/sediment;
- Incidental ingestion by wildlife for constituents in creek soil/sediment;
- Ingestion (drinking) by wildlife for constituents in creek surface waters; and
- Ingestion by wildlife for constituents that have bioaccumulated into riparian plants and benthic macroinvertebrate prey.

A conceptual site model (CSM) was prepared that identifies and summarizes the sources, mechanisms of transport, media of concern, exposure routes, and receptor groups and is intended to identify those exposure scenarios that are most likely to put BROCs at risk. A CSM for the Lower Red Butte Creek ERA is shown in Figure 5-1.

Bank soils are considered to be alluvial deposition of upstream sediments as a result of past high flow events. Although available to riparian wildlife, these sediments are not typically available to and were not quantitatively assessed for in-creek biota. For riparian wildlife (birds and mammals), creek soil/sediment (= bank soil + in-creek sediment) were evaluated.

***Inhalation of VOCs and Dermal Contact.*** VOC vapors are rapidly dispersed in aboveground air following volatilization from soil or surface

water. This dispersion, caused by wind and advection, results in very low exposure point concentrations of VOCs in aboveground air (USEPA 1998). Based on available information and previous experience, VOCs in outdoor air seldom “drive” risk (USEPA 2005). While potentially complete, inhalation exposure to VOCs is considered an insignificant exposure pathway for surface-dwelling wildlife (USEPA 2005).

Feathers of birds, fur on mammals, and scales on reptiles are believed to reduce dermal exposure by limiting the contact of the skin surface with the contaminated media (USEPA 2005). Accordingly, although potentially complete, dermal contact is considered an insignificant exposure pathway for wildlife (Peterle 1991; USEPA 2005).

Lack of a quantitative evaluation for these pathways is discussed in the uncertainty analysis (Section 8.6).

#### 5.4 ASSESSMENT ENDPOINTS

Assessment endpoints are “explicit expressions of the actual environmental value that is to be protected” (USEPA 1992a, 1998) and link the risk assessment to management concerns.<sup>19</sup> Assessment endpoints were established for this ERA to scope the risk assessment for in-creek aquatic biota and riparian wildlife BROCs (Table 5-3). Community-level assessment endpoints were established for aquatic and benthic macroinvertebrate communities. Population-level assessment endpoints were established for riparian wildlife (USEPA 1989).

**Table 5-3. Assessment Endpoints for Lower (Urban) Red Butte Creek**

Receptor	Level	Assessment Endpoint <sup>a</sup>
<b>In-Creek Aquatic Biota</b>		
Aquatic Plant	Community	Continued structural integrity of aquatic plant community
Aquatic Invertebrate	Community	Continued structural integrity of aquatic invertebrate community
Benthic	Community	Continued structural integrity of benthic

<sup>19</sup> Assessment endpoints are comprised of two elements: (1) the entity of concern and (2) a characteristic of the entity that is important to protect and is potentially at risk (USEPA 1992a, 1998).

Receptor	Level	Assessment Endpoint <sup>a</sup>
Macroinvertebrate		macroinvertebrate community
Fish	Population	Continued persistence of fish populations
Amphibian	Population	Continued persistence of amphibian populations
<b>Riparian Wildlife</b>		
Reptiles	Population	Continued persistence of reptile populations
Waterfowl/Shorebirds	Population	Continued persistence of waterfowl/shorebird populations
Mammals	Population	Continued persistence of riparian mammal populations
<i>Note:</i> a. Comparable to urbanized reaches of local background creeks in Salt Lake City.		

## 6.0 EXPOSURE ASSESSMENT

Exposure assessment establishes the information necessary to determine or predict ecological exposures to COPECs under exposure conditions of interest. Given the community coverage and/or home ranges of identified BROCs, the ERA evaluates the reach of Lower Red Butte Creek affected by the Spill<sup>20</sup> as a single exposure area. Exposures to wildlife receptors were estimated using exposure models consistent with USEPA's *Wildlife Exposure Factors Handbook* (1993).

### 6.1 TOTAL PETROLEUM HYDROCARBONS CONCENTRATIONS

It is noted that TPH in surface water and sediments were analyzed using USEPA Method 8015 (CPL 2011). This method does not report TPH in terms of carbon-chain fractions as needed when evaluating potential risks using the UDEQ approach. Accordingly and consistent with the HHRA, the results of the USEPA 8015 analyses were allocated to specific aliphatic/aromatic carbon-chain fractions using default (assumed) proportions provided by guidance (California Environmental Protection Agency [Cal/EPA] 2009) – i.e., 50-50 percent allocation between aromatic and aliphatic fractions.

### 6.2 EXPOSURE POINT CONCENTRATIONS

An exposure point concentration (EPC) is the concentration of a constituent in an environmental medium that a receptor of concern is likely to contact. In accordance with regulatory guidance, the lesser value of (i) the 95UCL on the mean or (ii) the maximum measured concentration in accessible media will be used to estimate exposure (USEPA 1989). All calculations of EPCs were performed using USEPA's ProUCL v. 4.01.00. To ensure consistency, McDaniel-Lambert conducted calculations of all EPCs for both the HHRA and this SLERA.

---

<sup>20</sup> From the Former Lower Underflow Dam (sampling location at the Spill site) to Below 900 East (furthest downstream sampling location).

### 6.3 *AQUATIC AND BENTHIC MACROINVERTEBRATES*

Exposures for in-creek aquatic and benthic macroinvertebrate biota will be reported in terms of concentrations in surface water and creek sediments, respectively.<sup>21</sup>

### 6.4 *RIPARIAN WILDLIFE*

In addition to environmental point concentrations, the essential inputs needed to estimate exposure to terrestrial wildlife are:

- Indicator species;
- Exposure equations;
- Wildlife exposure factors; and
- Biological uptake factors.

#### 6.4.1 *Indicator Species*

Given the number of species and the complexity of biological communities, all BROCs present in Lower Red Butte Creek cannot be individually assessed. Indicator species are identified to focus the ERA and evaluate risk for a representative set of species. Risks to indicator species are subsequently used to infer the potential for adverse impacts to taxonomically and functionally related BROCs. For further details on indicator species, please see Appendix A.

Indicator wildlife species for Lower Red Butte Creek ERA include:

- Mallard – herbivore (waterfowl);
- Spotted sandpiper – invertivore (shorebird);
- Musk rat – herbivore; and
- Raccoon – invertivore.

---

<sup>21</sup> Toxicity benchmarks for aquatic and benthic macroinvertebrate biota are in units of concentration for surface water and sediment, respectively (see Section 7).



To bound risk among herbivores and invertivores (invertebrate-consuming animals), indicator species were assumed to have a diet proportion of 100 percent for their particular food type (i.e., omnivores will be considered to have an exposure and risk intermediate to representative surrogate wildlife species).

#### 6.4.2 *Exposure Equations*

Exposures (or doses) are calculated using pathway-specific exposure equations for VOCs, SVOCs, and TPH. To facilitate comparisons with available toxicity data, estimates of exposure for COPECs will be reported in the units of dose,  $\text{mg}_{\text{COPEC}}/\text{kg}_{\text{body wt}}\text{-day}$ , using the following general equation (USEPA 1993):<sup>22</sup>

$$\text{Dose} = \text{EPC} \cdot \text{CR} \cdot \text{FC} \cdot \text{AF} \cdot \text{BW}^{-1} \quad \dots \text{Eq. 6-1}$$

where:

*EPC* = COPEC exposure point concentration for the medium of concern

*CR* = contact rate (e.g., ingestion rate)

*FC* = fraction of media contacted (e.g., diet proportions, proportion of time spent in Lower Red Butte Creek)

*AF* = assimilation factor<sup>23</sup>

*BW* = body weight

For specific exposure equations used in this SLERA, please see Appendix A.

#### 6.4.3 *Riparian Wildlife Exposure Factors*

In addition to EPCs, wildlife exposure factors are needed to evaluate exposure equations. To estimate exposures due to ingestion, the following wildlife exposure factors (WEFs) are required:

---

<sup>22</sup> Estimates of exposure to in-creek aquatic and benthic macroinvertebrate biota are in units of concentration, and, therefore, do not require exposure equations.

<sup>23</sup> Gut absorption factor is equal to the percent of concentration in surface water, sediment, or food that is absorbed across the gastrointestinal tract and is conservatively assumed to be 100 percent.

- Food ingestion and water (drinking) rates;
- Sediment and food diet proportions;
- Body weight; and
- Foraging area or home range.

In an effort to provide the most accurate assessment with the least amount of uncertainty, indicator species-specific data were used when available. When data for a selected indicator species were not available, data for a taxonomically related species having a similar feeding biology and size were used – if needed, metabolic adjustments were made. When no wildlife species-specific data were available, allometric regression equations provided in USEPA's *Wildlife Exposure Factor's Handbook* (1993) were used.

Wildlife exposure factors for the mallard, spotted sandpiper, muskrat, and raccoon are provided in Table 6-1.

**Table 6-1. Wildlife Exposure Factors**

Factor	Value	Source
<b>Mallard</b>		
Ingestion rate <sup>1</sup>	0.056 kg/kg-day	USEPA 1993
Drinking rate	0.0565 L/kg-day	USEPA 1993
Sediment diet proportion	3.3%	Beyer <i>et al.</i> 1994
Body weight <sup>2</sup>	1.134 kg	USEPA 1993
Home range <sup>3</sup>	580 ha	USEPA 1993
<b>Spotted Sandpiper</b>		
Ingestion rate <sup>1</sup>	0.163 kg/kg-day	USEPA 1993
Drinking rate	0.165 L/kg-day	USEPA 1993
Sediment diet proportion	8.2%	Beyer <i>et al.</i> 1994
Body weight <sup>4</sup>	0.052 kg	USEPA 1993
Home range	0.25 ha	USEPA 1993
<b>Muskrat</b>		
Ingestion rate <sup>5</sup>	0.30 kg/kg-day	USEPA 1993
Drinking rate	0.975 L/kg-day	USEPA 1993
Sediment diet proportion	9.4% <sup>a</sup>	Beyer <i>et al.</i> 1994
Body weight <sup>6</sup>	0.837 kg	USEPA 1993
Home range <sup>7</sup>	0.17 ha	USEPA 1993

Factor	Value	Source
<b>Raccoon</b>		
Ingestion rate <sup>1</sup>	0.537 kg/kg-day	USEPA 1993
Drinking rate	0.825 L/kg-day	USEPA 1993
Sediment diet proportion	9.4%	Beyer <i>et al.</i> 1994
Body weight <sup>8</sup>	3.99 kg	USEPA 1993
Home range <sup>9</sup>	156 ha	USEPA 1993
<i>Notes:</i> 1. Calculated from allometric equation (USEPA 1993). 2. Average of means from Nelson & Martin 1953, as cited in USEPA 1993. 3. Average of means from Kirby <i>et al.</i> 1985, as cited in USEPA 1993. 4. Average of means from Maxson & Oring 1980, as cited in USEPA 1993. 5. Average of means from Svihla & Svihla 1931, as cited in USEPA 1993. 6. Average of Reeves & Williams 1956, as cited in USEPA 1993. 7. Neal 1968, as cited in USEPA 1993. 8. Average of means from Johnson 1970, as cited in USEPA 1993. 9. Average of means from Stuewer 1943, as cited in USEPA 1993		

#### 6.4.4 *Biological Uptake Models*

For quantifying food chain exposures, simplified exposure models have been developed for terrestrial and aquatic food webs. COPEC concentrations transferred up the food chain will be calculated using available chemical-specific surface water-to-aquatic biota and sediment-to-benthic macroinvertebrate bioaccumulation factors (BAFs). BAFs used to calculate uptake into the prey of riparian wildlife are listed in Table 6-2 (also see Appendix E).

**Table 6-2. *Bioaccumulation Factors***

COPEC	BAFs		Source
	sed-to-plant <sup>1</sup>	sed-to-macroinvert <sup>2</sup>	
<b>TPH-Diesel / TPH-Motor Oil</b>			
Aromatics	1.2	1431	USEPA 2007
Aliphatics	0.54	17	USEPA 2007

• This page intentionally left blank •

## 7.0 EFFECTS ASSESSMENT

The effects assessment establishes toxicity reference values (TRVs) that are protective of aquatic biota, benthic macroinvertebrates, and wildlife. Ideally, the TRV is the highest dose or media concentration at which no chronic effects occur, and above which chronic adverse effects begin to occur.

Measures of effect are measurable responses to a stress that are related to and are used to evaluate the assessment endpoint (USEPA 1998). TRVs used in this ERA were used directly or derived from:

- Water quality criteria,
- Sediment quality guidelines, and
- Chronic reproductive or developmental impairment toxicity studies for birds and mammals.

ERM obtained TRVs that are protective of freshwater aquatic biota, benthic macroinvertebrate communities, and wildlife (birds, mammals) from widely recognized sources as summarized in Table 7-1 and discussed in the following sections.

**Table 7-1. Sources of Toxicity Reference Values**

TRVs	Surface Water	Sediment	Wildlife
Preferred	UT WQS MaDEP WQS	TEC/PEC MaDEP SQG	USEPA 2007
Alternative 1	NAWQC	NOAA SQuiRT	Sample et al. 1998
Alternative 2	Tier II WQS	Jones et al. 1997	TPHCWG 1997b
Alternative 3	USEPA EcoTox		
<i>Notes:</i> EcoTox =- USEPA Ecotox database NAWQC = National Ambient Water Quality Criteria TEC/PEC = Threshold effect concentration/probable effect concentration WQS = water quality standard			

## 7.1 **AQUATIC BIOTA TOXICITY REFERENCE VALUES**

For this SLERA, the following surface water benchmarks were used as TRVs for aquatic biota (in order of preference):

- State of Utah numerical water quality standards;
- MaDEP TPH numerical surface water guidelines (MaDEP 2002);
- National Recommended Water Quality Criteria (NRWQC) (USEPA 2009); and
- Tier II values (USEPA 1993, as cited in Suter & Tsao 1996).

No petroleum-related COPECs were identified for surface water in Lower Red Butte Creek. The lack of COPECs suggests that the Spill poses no potential risks to aquatic biota (aquatic plants, aquatic invertebrates, fishes).

**Table 7-2. Aquatic Biota Toxicity Reference Values**

COPEC	TRV (µg/L)	Source
None <sup>1</sup>		
Notes: 1 No COPECs were identified for surface water.		

## 7.2 **BENTHIC MACROINVERTEBRATE TOXICITY REFERENCE VALUES**

For this SLERA, the following sediment benchmarks were used as TRVs for aquatic biota (in order of preference):

- Threshold effect concentration (TEC)/probable effect concentrations (PEC) (MacDonald et al. 2000);
- MaDEP TPH sediment benchmarks (MaDEP 2007);
- USEPA Assessment and Remediation Contaminated Sediments (ARCS) Program values; and
- Oak Ridge National Laboratory (ORNL) toxicological benchmarks for sediment-associated biota (Jones et al. 1997).

TRVs for COPECs in sediment are listed in Table 7-3. TECs and PECs were used to bound potential risks as follows (McDonald et al. 2000):

- Less than TEC: adverse effects are not expected to occur;

- Between TEC and PEC: neither predicted to be toxic nor non-toxic – no guidance provided for this range; and
- Greater than PEC: adverse effects are expected to occur more often than not.

Note that MaDEP sediment quality guidelines for the protection of sediment-dwelling biota were used to evaluate potential risks due to residual exposures to aromatic and aliphatic fractions of TPH.

**Table 7-3. Benthic Macroinvertebrate Community Toxicity Reference Values**

COPEC	TRV (mg/kg)		Source
	TEC	PEC	
<b>Low Molecular Weight PAH</b>			
Anthracene	0.057	0.85	1
<b>High Molecular Weight PAH</b>			
Benzo(a)anthracene	0.11	1.1	2
Benzo(a)pyrene	0.15	1.5	2
Benzo(b)fluoranthene	0.11	1.1	2
Benzo(g,h,i)perylene	0.11	1.1	2
Benzo(k)fluoranthracene	0.11	1.1	2
Indeno(1,2,3-cd)pyrene	0.11	1.1	2
Pyrene	0.11	1.1	2
<b>TPH</b>			
<b>TPH Diesel</b>			
Aromatic	0.090	–	3
Aliphatic	3.2	–	3
<b>TPH Motor Oil</b>			
Aromatic	–	–	3
Aliphatic	9.9	–	3
<i>Notes:</i> 1 Dutch target and intervention values. 2 McDonald et al. (2000). 3 MaDEP (2007) – the lowest values for TPH-diesel and TPH- motor oil aromatic carbon-chains and aliphatic carbon-chains were used (consistent with HHRA).			

### 7.3

#### **RIPARIAN WILDLIFE TOXICITY REFERENCE VALUES**

For this SLERA, the following toxicity benchmarks were used as TRVs for wildlife (in order of preference):

- USEPA ecological soil screening levels (USEPA 2003-2007);

- ORNL toxicological benchmarks for wildlife (Sample et al. 1998); and
- TPHCWG toxicity benchmarks.

Wildlife TRVs for COPECs in creek soil/sediment are listed in Table 7-4.

**Table 7-4. Wildlife Toxicity Reference Values**

COPEC	TRV (mg/kg-day)		Source
	Bird	Mammal	
TPH			
TPH Diesel			
Aromatic	na	3.0	2
Aliphatic	na	10	2
TPH Motor Oil			
Aromatic	na	na	
Aliphatic	na	60	2
<i>Notes:</i> na = no TRV available 1. Sample et al. (1998). 2. TPHCWG (1997b) – the lowest values for TPH- diesel and TPH- motor oil aromatic carbon-chains and aliphatic carbon-chains were used (consistent with HHRA). The uncertainty factors for animal-to-human extrapolation and human variability were not included.			

#### 7.4 IN-CREEK BENTHIC MACROINVERTEBRATE COMMUNITY STRUCTURE

Findings of the benthic macroinvertebrate survey provide an additional line of evidence to characterize potential risks to in-creek biota. Quantitatively assessing the in-creek benthic macroinvertebrate community provides a number of advantages because they (USEPA 1999b):

- Indicate localized conditions given their limited migration patterns or sessile mode of life;
- Integrate effects of short-term environmental variations; and
- Constitute a broad range of trophic levels and pollution tolerances, thus providing strong information for interpreting cumulative effects.

The benthic macroinvertebrate community can be characterized in a number of ways, including measuring its diversity, its community composition, and its tolerance to perturbation. Specific metrics that allow



us to characterize the structure of the benthic macroinvertebrate community are listed in Table 7-5.

**Table 7-5. In-Creek Benthic Macroinvertebrate Community Metrics**

Category	Metric	Definition	Purpose
Diversity measures	Richness	Total number of taxa	Measures the overall variety of the macroinvertebrate assemblage
	Evenness	Relative abundance with which each species is represented in an area	Index of how close in numbers the species in the community are
	Diversity (Simpson's or Shannon-Wiener indices)	An index for the combined richness and evenness of species in the community	Provides a measure for the number of species weighted by their abundance
Composition measures	% EPT	Percent of composite of mayfly, stonefly, and caddisfly larvae	Measures the composite abundance of sensitive taxa; generally decreases after perturbation
	% Chironomidae	Percent of midge larvae	Measures the abundance of a tolerant taxon; generally increases after perturbation
Tolerance measures	Hilsenhoff biotic index (HBI)	Uses tolerance values to weight abundance in an estimate of overall pollution. Ranges from 1 to 10 (1 = pollutant sensitive to 10 = pollutant tolerant)	Perturbation should increase this value

• This page intentionally left blank •

## 8.0 RISK CHARACTERIZATION

Risk characterization integrates the results of the analysis phase (i.e., exposure and effects assessments) to evaluate the likelihood of adverse ecological impacts associated with exposure to COPECs (USEPA 1992a). Potential risks were inferred based on:

- Hazard quotients (HQs); and
- Analysis of the benthic macroinvertebrate community structure.

Potential risks were evaluated for:

- Aquatic biota community;
- Benthic macroinvertebrate community;
- Riparian bird populations; and
- Riparian mammal populations.

## 8.1 HAZARD QUOTIENTS

HQs are used to estimate the potential for adverse ecological impacts when sufficient exposure and toxicity data exist. An HQ is simply the ratio of the estimated exposure to the TRV:

$$\text{HQ} = \frac{\text{Estimated Exposure}}{\text{TRV}}$$

An HQ less than one ( $\text{HQ} < 1$ ) indicates a negligible potential for adverse ecological impacts due to exposure to a particular COPEC, whereas an HQ greater than or equal to one ( $\text{HQ} \geq 1$ ) indicates a potential for adverse ecological impacts due to exposure to that COPEC.

The hazard index (HI) is the sum of HQs ( $\text{HI} = \sum \text{HQs}$ ) and was calculated to evaluate potential cumulative risks for constituents with similar structure activity relationships. Similar to HQs, an HI less than one ( $\text{HI} < 1$ ) indicates a negligible potential for adverse ecological impacts due to cumulative exposures to COPECs, whereas an HI greater than or equal to one ( $\text{HI} \geq 1$ ) indicates a potential for adverse ecological impacts due to cumulative exposures to COPECs.

Risk estimates (HQs and HIs) for aquatic biota, benthic macroinvertebrates, riparian birds, and riparian mammals are provided in Tables 8-1 through 8-6 (see end of report; also see Appendix E).

## 8.2 *AQUATIC BIOTA*

Aquatic biota includes aquatic plants, aquatic (pelagic water column) invertebrates, and fishes (Section 5.1). Numerical water quality criteria used to evaluate potential risks in this SLERA were established by regulatory agencies to be protective of aquatic biota communities.

In Lower Red Butte Creek, all VOCs, PAHs, and TPH were either detected in less than 5 percent of surface water samples or had maximum detected concentrations less than risk-based ESLs.

The only constituent identified for surface waters of Lower Red Butte Creek was the SVOC bis(2-ethylhexyl)phthalate. This phthalate ester was detected in only 1 of 16 surface water samples (6 percent frequency of detection). Further, bis(2-ethylhexyl)phthalate is not considered a petroleum-related constituent (i.e., not related to a crude oil release) and is not evaluated further in this SLERA.

## 8.3 *BENTHIC MACROINVERTEBRATES*

In Lower Red Butte Creek, nearly all VOCs were either detected in less than 5 percent of creek bed sediment samples or had maximum detected concentrations less than risk-based ESLs. The only VOCs detected in creek bed sediments of Lower Red Butte Creek were acetone, PCE, and toluene, each of which was detected in only 1 of 16 creek bed samples. Note that acetone and PCE are not considered petroleum-related constituents (i.e., not related to a crude oil release); further, acetone is a common laboratory contaminant. Toluene was not detected at a concentration greater than the ESL for creek sediments. Hence, acetone, PCE, and toluene are not evaluated further in this SLERA.

Other COPECs include eight PAHs and TPH-diesel and TPH-motor oil. The COPECs in creek bed sediments of Lower Red Butte Creek are summarized in Table 8-2.

### 8.3.1 Hazard Quotients

Sediment quality guidelines used to evaluate potential risks in this SLERA were established by regulatory agencies to be protective of sediment-dwelling benthic communities.

EPCs for anthracene and indeno(1,2,3-cd)pyrene<sup>24</sup> in Lower Red Butte Creek bed sediments resulted in HQs of 1.6 and 1.1, respectively. The EPCs for anthracene and indeno(1,2,3-cd)pyrene do not exceed the PEC sediment quality guideline – hence, these COPECs may or may not pose a risk to benthic macroinvertebrate communities in Lower Red Butte Creek.

TPH-diesel and TPH-motor oil were frequently detected in creek bed sediments of Lower Red Butte Creek. EPCs for aromatic and aliphatic fractions of TPH results in HQs greater than one (HQs > 1), suggesting that these fractions pose a potential risk to benthic macroinvertebrate communities. However, as noted in Section 4, concentrations of TPH-diesel and TPH-motor oil measured in creek bed sediment of Lower Red Butte Creek were comparable to or less than concentrations measured in creek bed sediment of local background creeks (Table 8-2a,b), as shown below:

COPEC	Lower Red Butte Creek	Local Background Creeks
<b>TPH-Diesel</b>		
Aromatics	49	77
Aliphatics	49	77
<b>TPH-Motor Oil</b>		
Aromatics	53	49
Aliphatics	53	49
* Values are concentrations (mg/kg)		

These EPCs suggest that residual exposures/risks of TPH in creek bed sediments of Lower Red Butte Creek may be comparable to local

<sup>24</sup> Anthracene was detected in only 2 of 15 sampling locations; while, indeno(1,2,3-cd)pyrene was detected in 3 of 15 samples.

background creeks. Uncertainties related to potential risk estimates are discussed further in Section 8.6.

### 8.3.2 *Community Structure Metrics*

Benthic macroinvertebrates are intimately associated with in-creek bed sediments and assemblages of macroinvertebrates are considered good indicators of localized sediment quality conditions because (USEPA 1999b):

- Benthic macroinvertebrates have limited migration patterns or a sessile mode of life and are particularly well-suited for assessing site-specific impacts.<sup>25</sup>
- Benthic macroinvertebrates integrate the effects of short-term environmental variations.<sup>26</sup>
- Benthic macroinvertebrate assemblages are made up of species that constitute a broad range of trophic levels and pollution tolerances.
- Benthic macroinvertebrates provide strong information for interpreting cumulative effects.

Benthic macroinvertebrates were collected (Table 8-7), identified, and enumerated to provide additional lines of evidence to support the SLERA and are intended to assist in characterizing/verifying potential impacts to benthic macroinvertebrate communities that may be attributable to exposures to in-creek sediments.

**Table 8-7. *Benthic Macroinvertebrate Sampling Locations within Lower Red Butte Creek***

<b>Sampling Site</b>	<b>Description</b>	<b>Distance (miles)</b>
Red Butte Gardens	Natural	0
Below Chipeta Way	Urban	0.31
Above Foothill Drive	Urban	0.85
At Mt Olivet Diversion	Urban	1.2

<sup>25</sup> Also upstream/downstream comparisons.

<sup>26</sup> Most species have a complex life cycle of approximately 1 year or more.

Sampling Site	Description	Distance (miles)
Above Sunnyside Avenue	Urban	1.3
At County Stream Gauging Station	Urban	1.7
Above 1500 East	Urban	1.9
Below 1300 East	Urban	2.3
Below 1100 East	Urban	2.6

For further details on the findings of these analyses, please see Appendix F.

### 8.3.2.1 *Comparison to Local Background Creeks*

Nearby urban creeks were selected to represent in-creek local background conditions having similar biological expectations as Lower Red Butte Creek in the absence of the effects of the Spill. As such, local background creeks can be used to characterize the "reasonably attainable" state and can provide the point-of-reference to assess the potential impairment as a result of the Spill.

Boxplots (Figure 8-1) were constructed and nonparametric Wilcoxon Rank Sum (WRS) two-sample tests were conducted on the following community metrics:

- Taxa richness
- Evenness
- Shannon's diversity index
- Hilsenhoff biotic index (HBI)
- Percent chironomids
- Percent EPT

For each metric, the WRS tests evaluated the hypothesis that:

$H_0$ : There is no significant difference in median metric values between Lower Red Butte Creek and local background creek.

Both boxplots and WRS<sup>27</sup> tests suggest that there are no differences between Lower Red Butte Creek and local background creeks for each/all of the community metrics (Table 8-8).

**Table 8-8. Benthic Macroinvertebrate Community: Results of Nonparametric Two-Sample Tests**

H<sub>0</sub>: No significant difference between Lower Red Butte Creek and local background creeks

Metric	<i>p</i>	Significant?
Richness	0.13	<i>ns</i>
Evenness	0.42	<i>ns</i>
Shannon-Wiener Diversity Index	0.25	<i>ns</i>
% EPT	0.38	<i>ns</i>
% Chironomids	0.56	<i>ns</i>
Hilsenhoff biotic index (HBI)	0.59	<i>ns</i>
Notes: <i>ns</i> = Not significant <i>p</i> = Probability		

These findings are consistent with findings of the Local Background Evaluation (Section 4) that there is no difference in PAH concentrations in creek sediments between Lower Red Butte Creek and the local background creeks.

### 8.3.2.2 Trends in Community Metrics Relative to Distance from Spill

To test the hypothesis that impacts to benthic macroinvertebrates would be greatest nearest the Spill and decrease with distance downcreek, a regression analysis was conducted for the same set of metrics as listed in Section 8.3.2.1. For each metric, the regression evaluated the hypothesis that:

H<sub>0</sub>: There is no significant trend with distance from the Spill site downstream to lower portions of Lower Red Butte Creek.

---

<sup>27</sup> At a significance level ( $\alpha$ ) of 0.05, results of WRS tests show none of the community metrics are significantly different between Lower Red Butte Creek and the local background creeks.



The regression analyses found that (Table 8-9):

- Taxa richness and percent EPT (pollutant sensitive) decreased with distance downstream from the Spill site;
- Percent chironomids (pollutant tolerant) increased with distance downstream from the Spill site;
- Evenness, diversity, and HBI<sup>28</sup> demonstrated no significant changes with distance downstream from the Spill site; and
- Benthic macroinvertebrate metrics from samples collected at locations immediately downstream from the Spill site were “healthier” (i.e., higher richness, higher percent EPT, lower percent chironomids) than samples collected farther downstream in the more urbanized reach of Lower Red Butte Creek.

**Table 8-9. Benthic Macroinvertebrate Community: Results of Nonparametric Regression Analyses**

H<sub>0</sub>: No significant trend with distance downstream of Spill site in Lower Red Butte Creek.

Metric	Sign of Slope	p	Significant?
Richness	negative	0.05	*
Evenness	negative	0.72	ns
Shannon-Wiener Diversity Index	negative	0.16	ns
% EPT	negative	0.08	ns
% Chironomids	positive	0.05	*
Hilsenhoff biotic index (HBI)	positive	0.25	ns
Notes: ns = Not significant * = Significant at significance level ( $\alpha$ ) = 0.05			

These findings suggest that impacts to benthic macroinvertebrates increase with distance downstream (further away) from the Spill site. Two possible conclusions include impacts are due to (i) Spill-related hydrocarbons that were transported and deposited in downstream locations or (ii) urban runoff-related hydrocarbons deposited in lower

<sup>28</sup> Note that HBI ranges from 1 (pollutant sensitive) to 10 (pollutant tolerant) – i.e., higher values indicate community composed of more pollutant-tolerant taxa.

creek locations. The Local Background Evaluation found that the PAH composition in Lower Red Butte Creek appear to be consistent with sources typical of urban runoff (Section 4). These findings suggest that the source of downstream hydrocarbons is likely from urban runoff sources.

## 8.4 *RIPARIAN BIRDS*

Riparian birds were considered to be exposed to COPECs in surface waters and creek soil/sediments (Section 5.3). In Lower Red Butte Creek, all VOCs, SVOCs, and PAHs were either detected in less than 5 percent of creek soil/sediment samples or had maximum detected concentrations less than risk-based ESLs. No COPECs were identified for surface waters and TPH-diesel and TPH-motor oil were the only COPECs identified for creek soils/sediments of Lower Red Butte Creek.

### 8.4.1 *Mallard*

Risk estimates for the mallard were used to infer potential risks to birds consuming riparian emergent plants (herbivores).

No avian TRVs were identified for aromatic and aliphatic fractions of TPH-diesel and TPH-motor oil. Hence, no risk estimates were quantified for TPH in creek soil/sediment for these COPECs (Table 8-3).

Note that nearly 95 percent of the exposure to aromatic and 85 percent of the exposure to aliphatic fractions of TPH is due to the ingestion of plants. Doses due to ingestion were derived using  $k_{ow}$ -modeled BAFs. These BAFs suggest biomagnification of TPH in plants and may not be representative of tissue burdens of plants in Lower Red Butte Creek (also see Section 8.6).

Moreover, as noted in Section 4, concentrations of TPH-diesel and TPH-motor oil measured in soil/sediment of Lower Red Butte Creek were comparable to or less than concentrations measured in soil/sediment of local background creeks. As shown below, exposures (doses)<sup>29</sup> to aliphatic and aromatic fractions of TPH-diesel are greater in local

---

<sup>29</sup> Given the lack of avian TRVs, exposures (doses) provide a means to compare Lower Red Butte Creek and local background creeks.

background creeks as compared to Lower Red Butte Creek (from Table 8-3a,b):

COPEC	Lower Red Butte Creek	Local Background Creeks
<b>TPH-Diesel</b>		
Aromatics	3.6	3.7
Aliphatics	1.6	1.7
<b>TPH-Motor Oil</b>		
Aromatics	3.4	2.3
Aliphatics	1.5	1.1
* Values are doses (mg/kg-day)		

These data suggest that residual exposures/risks of TPH in creek soil/sediments of Lower Red Butte Creek may be comparable to local background creeks.

Uncertainties related to potential risk estimates are discussed further in Section 8.6.

#### 8.4.2 *Sandpiper*

Risk estimates for the sandpiper were used to infer potential risks to birds consuming benthic macroinvertebrates (invertivores).

No avian TRVs were identified for aromatic and aliphatic fractions of TPH-diesel and TPH-motor oil. Hence, no risk estimates were quantified for TPH in creek soil/sediment for these COPECs (Table 8-4).

Note that exposures to aromatic and aliphatic fractions of TPH appear to be high. Nearly 100 percent of the exposure to aromatic and aliphatic fractions of TPH is due to the ingestion of macroinvertebrate. Doses due to ingestion were derived using  $k_{ow}$ -modeled BAFs (also see Section 8.6). BAFs for aromatic fractions are on the order of  $10^{+3}$ , while BAFs for aliphatic fractions are on the order of  $10^{+1}$ . These BAFs appear to be particularly high, suggesting biomagnification of TPH in benthic macroinvertebrates, and may not be representative of tissue burdens of benthic macroinvertebrates in Lower Red Butte Creek (also see Section 8.6).

Moreover, as noted in Section 4, concentrations of TPH-diesel and TPH-motor oil measured in soil/sediment of Lower Red Butte Creek were

comparable to or less than concentrations measured in soil/sediment of local background creeks. As shown below, exposures (doses) to aliphatic and aromatic fractions of TPH-diesel are greater in the local background creeks as compared to Red Butte Creek (from Table 8-4a,b):

COPEC	Lower Red Butte Creek	Local Background Creeks
<b>TPH-Diesel</b>		
Aromatics	11931	12258
Aliphatics	145	149
<b>TPH-Motor Oil</b>		
Aromatics	11181	7839
Aliphatics	136	95
* Values are doses (mg/kg-day)		

These data suggest that residual exposures/risks of TPH in creek soil/sediments of Lower Red Butte Creek may be comparable to the local background creeks.

Uncertainties related to potential risk estimates are discussed further in Section 8.6.

## 8.5 **RIPARIAN MAMMALS**

Riparian mammals were considered to be exposed to COPECs in surface waters and creek soil/sediments (Section 5.3). In Lower Red Butte Creek, all VOCs, SVOCs, and PAHs were either detected in less than 5 percent of creek soil/sediment samples or had maximum detected concentrations less than risk-based ESLs. No COPECs were identified for surface waters and TPH-diesel and TPH-motor oil were the only COPECs identified for creek soils/sediments of Lower Red Butte Creek.

### 8.5.1 **Muskrat**

Risk estimates for the muskrat were used to infer potential risks to mammals consuming riparian emergent plants (herbivores).

Risk estimates for the muskrat are provided in Table 8-5. HQs suggest that exposures to the aromatic fraction of TPH-diesel and cumulative exposures to TPH pose a potential risk (HQ > 1).

Note that nearly 95 percent of the exposure to aromatic and 85 percent of the exposure to aliphatic fractions of TPH is due to the ingestion of plants. Doses due to ingestion were derived using  $k_{ow}$ -modeled BAFs. These BAFs suggest biomagnification of TPH in plants and may not be representative of tissue burdens of plants in Lower Red Butte Creek (also see Section 8.6).

Moreover, as noted in Section 4, concentrations of TPH-diesel and TPH-motor oil measured in soil/sediment of Lower Red Butte Creek were comparable to or less than concentrations measured in soil/sediment of local background creeks. As shown below, exposures (doses<sup>30</sup>) to aliphatic and aromatic fractions of TPH-diesel are greater in local background creeks as compared to Red Butte Creek (from Table 8-5a,b):

COPEC	Lower Red Butte Creek	Local Background Creeks
<b>TPH-Diesel</b>		
Aromatics	20	21
Aliphatics	9.7	9.9
<b>TPH-motor oil</b>		
Aromatics	19	13
Aliphatics	9.1	6.4
* Values are doses (mg/kg-day)		

However, the HQ due to exposure to the aliphatic fraction<sup>31</sup> of TPH-motor was greater than one and exposures were greater in soil/sediments of Lower Red Butte Creek as compared to local background creeks.

Uncertainties related to potential risk estimates are discussed further in Section 8.6.

---

<sup>30</sup> Consistent with discussion for indicator birds, exposures (doses) are compared.

<sup>31</sup> The aromatic fraction of TPH-motor oil could not be quantitatively evaluated due to the lack of a mammalian TRV.

## 8.5.2

**Raccoon**

Risk estimates for the raccoon were used to infer potential risks to mammals consuming benthic macroinvertebrates (invertivores).

Risk estimates for the raccoon are provided in Table 8-6. HQs suggest that exposures to the aromatic fraction of TPH-diesel and cumulative exposures to TPH pose a potential risk (HQ >> 1).

Note that exposures to aromatic and aliphatic fractions of TPH appear to be high. Nearly 100 percent of the exposure to aromatic and aliphatic fractions of TPH is due to the ingestion of macroinvertebrate. Doses due to ingestion were derived using  $k_{ow}$ -modeled BAFs (also see Section 8.6). BAFs for aromatic fractions are on the order of  $10^{+3}$ , while BAFs extraordinarily aliphatic fractions are on the order of  $10^{+1}$ . These BAFs appear to be particularly high, suggest biomagnification of TPH in benthic macroinvertebrates, and may not be representative of tissue burdens of benthic macroinvertebrates in Lower Red Butte Creek (also see Section 8.6).

However, as noted in Section 4, concentrations of TPH-diesel and TPH-motor oil measured in soil/sediment of Lower Red Butte Creek were comparable to or less than concentrations measured in soil/sediment of local background creeks. As shown below, exposures (doses) to aliphatic and aromatic fractions of TPH-diesel are greater in local background creeks as compared to Red Butte Creek (from Table 8-6a,b):

COPEC	Lower Red Butte Creek	Local Background Creeks
<b>TPH-Diesel</b>		
Aromatics	39307	40383
Aliphatics	477	490
<b>TPH-Motor Oil</b>		
Aromatics	36836	25824
Aliphatics	447	313
* values are doses (mg/kg-day)		

However, the HQ due to exposure to the aliphatic fraction of TPH-motor was greater than one and exposures were greater in soil/sediments of Lower Red Butte Creek as compared to local background creeks.

Uncertainties related to potential risk estimates are discussed further in Section 8.6.

## 8.6 **UNCERTAINTY ANALYSIS**

Consistent with USEPA (1989) guidance, a qualitative discussion of the uncertainties associated with the estimation of risks for Lower Red Butte Creek will be presented in the ERA report. This uncertainty analysis discusses key uncertainties associated the exposure and effects assessments. To reduce uncertainties, focused verification of SLERA analyses may be considered.

### 8.6.1 ***Uncertainties Associated with the Exposure Assessment***

Sources of uncertainty related to COPEC exposures include:

- Constituents detected in less than 5 percent of the samples;
- Bis(2-ethylhexyl)phthalate;
- Use of indicator species;
- Bioaccumulation factors;
- Omission of inhalation and dermal contact; and
- Comparisons to local background creek conditions.

#### 8.6.1.1 ***Constituents Detected in Less Than 5 Percent of the Samples***

With the exception of soil/sediment, other media of concern had sample sizes less than 20 samples (i.e., surface water and creek bed sediments). Hence, a frequency of detection (FOD) less than 5 percent was equivalent to “not detected in any sample”. Consistent with standard practice, petroleum-related compounds that were never detected and having detection limits less than COPEC screening levels were not identified as COPECs in this SLERA.

2-Methylnaphthalene and fluorene were detected in 1 of 27 soil/sediment samples from Lower Red Butte Creek. Hence, these petroleum-related constituents in soil/sediment that were detected at a frequency greater than 0 percent, but less than 5 percent ( $0\% < \text{FOD} < 5\%$ ) include:

- 2-Methylnaphthalene
- Fluorene

Note that both of the above constituents would not be identified as COPECs, as each was detected at a concentration less than their COPEC screening levels (see also Appendix E, *Ecological Risk Analyses*).

#### 8.6.1.2 *Bis(2-ethylhexyl)phthalate, Acetone, and Tetrachloroethene (PCE)*

Bis(2-ethylhexyl)phthalate was detected in 1 of 16 surface water samples from Lower Red Butte Creek; acetone and PCE were detected in 1 of 16 sediment samples from Lower Red Butte Creek. However, bis(2-ethylhexyl)phthalate, acetone, and PCE are not considered petroleum-related constituents – i.e., are not related to a crude oil release. These constituents were not evaluated further in this SLERA (as consistent with the HHRA).

#### 8.6.1.3 *Indicator Species*

Given the number of species and the complexity of biological communities, all species present in Lower Red Butte Creek cannot be individually assessed. Indicator species were used to infer the potential for adverse impacts to taxonomically and functionally related BROCs. To minimize the chance of underestimating exposure, indicator species were selected to maximize estimates of exposure (e.g., small body size, small home or foraging ranges, forages on ground surface), where possible. However, very little is known about the relative sensitivity to petroleum-related constituents among related species. Therefore, the extrapolation of risks from species to species introduces an unquantifiable amount of uncertainty.

#### 8.6.1.4 *Inhalation and Dermal Exposure Pathways*

VOC vapors are rapidly dispersed in aboveground air following volatilization from soil or surface water and are considered to result in very low exposure point concentrations of VOCs in aboveground air (USEPA 1998). While potentially complete, inhalation exposure to VOCs is considered an insignificant exposure pathway for surface-dwelling wildlife (USEPA 2005).

Feathers of birds, fur on mammals, and scales on reptiles are believed to reduce dermal exposure by limiting the contact of the skin surface with the contaminated media (USEPA 2005). Although potentially complete, dermal contact is considered an insignificant exposure pathway for wildlife (Peterle 1991; USEPA 2005).



Hence, quantitative evaluation of inhalation and dermal exposure pathways were omitted from this SLERA. Omission of inhalation and dermal contact exposure pathways may result in an underestimate of potential risks.

#### 8.6.1.5 *Bioaccumulation Factors*

BAFs for aromatic and aliphatic fractions of TPH-diesel and TPH-motor oil were derived from  $K_{ow}$  models (USEPA 2007). These  $K_{ow}$  models appear to have resulted in inordinately high BAFs for TPH – in particular, for sediment-to-benthic invertebrate BAFs for the aromatic fraction of TPH-diesel and TPH-motor oil.  $K_{ow}$ -modeled BAFs introduce an unquantifiable uncertainty into calculations of risk estimates for this SLERA. Consequently, exposures and risk estimates for aromatic and aliphatic fractions of TPH are considered to be uncertain.

To reduce uncertainty in the uptake, exposure, and potential risks via the foodchain, modeled BAFs may be verified by direct measures of tissue burdens of plants and benthic macroinvertebrates in the field. Given that exposures and potential risks to TPH in Lower Red Butte Creek are comparable or less than in local background creeks, verification of BAFs may not be considered critical to support decision-making.

#### 8.6.1.6 *Comparisons to Local Background Conditions*

For the purposes of this SLERA, comparisons to local background creek concentrations were not used to exclude/eliminate constituents from further evaluation in the SLERA. All constituents with greater than 5 percent frequency of detect and maximum concentrations greater than risk-based ESLs were evaluated further.

Visual examination of Q-Q plots indicated inconsistencies with statistical tests for some PAHs and TPH-motor oil. However, the Local Background Evaluation suggests that, for the most part, PAHs and TPH-diesel are consistent with local background creek conditions (McDaniel-Lambert 2012). Further, the PAH composition analysis suggests that petroleum-related hydrocarbons detected in Red Butte Creek appear to be consistent with PAHs typical of urban run-off. Hence, findings of the Local Background Evaluation suggest that petroleum hydrocarbons detected in Red Butte Creek may not be Spill-related.

## 8.6.2 *Uncertainties Associated with the Effects Assessment*

Sources of uncertainty related to effects include:

- Use of chronic no observable adverse effect level (NOAEL)-equivalent TRVs;
- Species-to-species toxicity extrapolations;
- Laboratory-to-field toxicity extrapolations;
- Constituent-to-constituent extrapolations; and
- Lack of avian TRVs for TPH.

### 8.6.2.1 *Use of Chronic No Observable Adverse Effect Level-Equivalent Toxicity Reference Values*

The use of chronic NOAEL-equivalent TRVs is likely to result in conservative assessments of risk because environmental exposures were compared to toxicity levels at which no adverse effects were observed. Studies indicate that acute LD<sub>50</sub>s (lethal dose for 50 percent of the population) derived from multiple dose toxicity tests show a high positive correlation with observed impacts in the environment (USEPA 1991). Cal/EPA (1996) considers chronic NOAELs to be 500 times more sensitive than LD<sub>50</sub>s and 10 times more sensitive than acute lowest observed adverse effect levels (LOAELs). Thus, use of chronic NOAEL-equivalent TRVs provides a substantially greater level of protection than the use of LOAELs or LD<sub>50</sub>s.

### 8.6.2.2 *Species-to-Species Toxicity Extrapolations*

A source of uncertainty in the SLERA is the lack of applicable species-specific toxicity data. Because of this data limitation, TRVs were developed using available toxicity data for laboratory test species. For example, TRV for the muskrat and raccoon were developed from toxicity data for mice and rats.

Species vary with respect to sensitivity to specific chemicals (Calabrese and Baldwin 1993). Based on our review of the toxicological data, the range of sensitivity for members within a class of vertebrates were typically up to 100-fold. This range of uncertainty is substantiated by Calabrese and Baldwin (1993). Although a range in sensitivity may be described, the relative sensitivity (and the “direction” of sensitivity) of riparian wildlife species compared to laboratory test species to COPEC exposures is not known.

### 8.6.2.3 *Laboratory-to-Field Toxicity Extrapolations*

A number of studies (primarily for aquatic systems) have evaluated the ability of single-chemical laboratory toxicity test results to predict adverse effects of that chemical on organisms under field conditions. Preliminary chemical contaminant studies suggest that laboratory toxicity tests represent more conservative exposure scenarios than those that occur in nature (USEPA 1991). Furthermore, concentrations of chemicals causing no effect in laboratory tests also do not appear to affect communities in the field. Thus, the use of chronic NOAEL-equivalent TRVs are likely to provide a conservative level of protection to plant and wildlife communities and populations observed in the field.

### 8.6.2.4 *Constituent-to-Constituent Extrapolations*

Sufficient toxicity data to develop reference toxicity values for all constituents was not possible. To assess risks, constituent-to-constituent extrapolations were required. Use of constituent-to-constituent extrapolations is supported by the abundance of research work on quantitative structure-activity relationships (QSAR) reported in the pharmaceutical and medicinal chemistry literature that suggests that chemicals with similar molecular or physicochemical properties have similar biological reactivity and toxicity (Donkin 1994; Nirmalakhandan and Speece 1988). The use of constituent-to-constituent extrapolations is also consistent with guidance for HHRA.

### 8.6.2.5 *Lack of Avian Toxicity Reference Values for Total Petroleum Hydrocarbons and Polycyclic Aromatic Hydrocarbons*

Hazardous components of TPH include PAHs and BTEX.<sup>32</sup> The lack of avian ecotoxicity data for TPH and PAHs hampers the quantitative risk assessment of risk for birds. This uncertainty is not unique to this SLERA.

---

<sup>32</sup> Benzene and ethylbenzene were not detected in creek soils/sediments. Although detected in 1 of 16 sediment samples, toluene was not detected at a concentration greater than the ESL for creek sediments. BTEX analytical detection limits are nearly two orders of magnitude less than these ESLs (see Table E-3a).

TPH and several PAHs were detected in creek soils/sediments (see Table E-3a of Appendix E). However, there are no widely recognized/accepted avian TRVs for TPH and PAHs. PAHs detected in creek soils/sediments were not present at concentrations greater than USEPA 2007 ecological soil screening levels (ESSLs) for PAHs that are protective of wildlife. However, USEPA ESSLs for wildlife were based on mammalian screening levels – avian ecotoxicity data were not considered sufficient to develop TRVs.

Because TPH and PAHs are of interest for the Spill, in lieu of no assessment, one may consider a tempered, qualitative comparison to risk estimates for mammals as a point-of-reference. For example, the aliphatic fraction of TPH-motor oil posed a risk to the muskrat and raccoon.<sup>33</sup> If birds are as sensitive to the aliphatic fraction of TPH-motor oil as small mammals, a similar pattern in potential risk would be observed. Note that the relative sensitivities between birds and mammals to the aliphatic fraction of TPH-motor oil is not known.

The lack of avian TRVs for TPH and PAHs is a data gap shared among any current toxicity-based assessment of risk for birds exposed to petroleum-related constituents and is considered a fundamental uncertainty in this SLERA.

### 8.6.3 *Comparison of Reach-Wide to Location-Specific Estimates of Risk*

Given the lack of state and federal listed species, assessment endpoints for the Lower Red Butte Creek identified BROCs at the population- and community-level (see Section 5.4). Hence, Lower Red Butte Creek was evaluated as a single exposure area/reach – a reach-wide average was considered the appropriate spatial scale to evaluate exposures and potential risks to populations and communities.

To summarize Sections 8.3 and 8.5, potential risks to benthic macroinvertebrates and riparian mammals due to residual reach-wide exposures to aromatic and aliphatic fractions of TPH in Lower Red Butte Creek were identified. Note that exposures to aromatic and aliphatic

---

<sup>33</sup> Note that the HQs for aromatic and aliphatic fractions of TPH-diesel were less than the corresponding HQs for local background creek.

fractions of TPH-diesel were comparable to or less than those measured/calculated for the local background creeks. Whereas, risk estimates for the aliphatic fraction of TPH-motor oil were greater than one and greater than risk estimates calculated for the local background creeks. The aromatic fraction of TPH-motor oil could not be quantitatively evaluated due to the lack of a mammalian TRV.

In response to a comment from UDEQ, location-specific risk estimates (HIs) were calculated, are provided in Appendix H, summarized in Table 8-10, and mapped in Figure 8-2. In general, location-specific risk estimates corroborate the findings of the reach-wide assessment (see Tables H-4a-c). However, location-specific risk estimates also appear to suggest that risks to benthic macroinvertebrates and muskrat (idealized herbivorous riparian mammal) may be localized (Table 8-10; Figure 8-2) – whereas, potential risks to the raccoon (idealized invertivore riparian mammal) may be reach-wide.

**Table 8-10. Summary of Location-Specific Risk Estimates for TPH**

Location	Macroinvertebrate	Muskrat	Raccoon
<b>Avg. Local Background</b>	190	12	13515
Underflow Dam	70	5.4	9156
Below Chipeta	208	8.9	14669
University Marriott	210	8.9	14919
Above Foothill	181	7.6	12860
Mt Olivet	129	5.5	9261
Above Sunnyside	109	10.8	18135
1731 E900	160	10.5	18262
Gaging Station	207	9.3	15690
Above 1500E	225	9.9	16595
Below 1300E	110	12.5	21218
Below 1100E	100	4.3	7332
Below 900E	92	3.9	6558

Values are TPH HIs

Highlighted cells are greater than HIs for upper bound average exposures in local background urban creeks (from Tables 8-1b to 8-6b)

Macroinvertebrate = benthic macroinvertebrates

The HHRA noted that the small sediment sample size at each creek location remains "...a significant uncertainty..." and could result in either an over- or underestimate of location-specific exposures/risks. For population- and community-level and/or ranging biotic receptors of

concern, this SLERA concurs with the HHRA in concluding that the creek-wide risk estimates are a more appropriate and reliable measures of potential exposure to petroleum-related constituents.

Further, location-specific risk estimates should be used with care if supporting decision-making related to the possible need and scope of location-specific clean-up and/or monitoring. For example, location-specific concentrations of TPH may be due to location-specific anthropogenic contributions that are unrelated to the spill. Given the original scope of this SLERA, no survey or evaluation of location-specific non-spill-related sources was performed to support characterizations of potential location-specific risks to biota.

## 9.0 DISCUSSION AND CONCLUSIONS

A key feature of this SLERA is the use of multiple lines of evidence (where available) to support characterizations of risk. The use of multiple lines of evidence (e.g., local background creek evaluation, risk estimates for local background creeks, benthic macroinvertebrate community metrics) is intended to provide several perspectives to assist in characterizing the potential for ecological risk.

**Aquatic Biota.** Potential risks to aquatic biota due to residual exposures of petroleum-related constituents in surface water appear to be negligible. All analytes, except for bis(2-ethylhexyl)phthalate, were either not detected or had maximum concentrations less than risk-based ESLs. Note that bis(2-ethylhexyl)phthalate was only detected in 1 of 16 surface water samples and this compound is not considered a petroleum-related constituent .

**Benthic Macroinvertebrate Community.** Potential risks to benthic macroinvertebrate communities due to residual exposures of petroleum-related constituents in creek bed sediments also appear to be limited. COPECs were limited to eight PAHs, TPH-diesel, and TPH-motor oil. Exposures for 6 of 8 PAHs were less than TECs leading to a conclusion that adverse effects are not expected to occur (MacDonald et al. 2000). Exposures to anthracene were greater than TECs, but less than PECs, and a determination of toxicity or nontoxicity cannot be confidently predicted (MacDonald et al. 2000).

An evaluation of local background creeks found that concentrations of PAHs in creek bed sediments of Lower Red Butte Creek were comparable or less than concentrations in creek bed sediments of local background creeks. In addition, exposures used in this SLERA appear to be comparable between Lower Red Butte Creek and local background creeks. Finally, metrics suggest that the structure<sup>34</sup> of the benthic macroinvertebrate community in Lower Red Butte Creek is comparable to local background creeks. These lines of evidence suggest that potential

---

<sup>34</sup> As characterized by richness, diversity, evenness, percent EPT, percent chironomids, and HBI.

exposures/risks in Lower Red Butte Creek are comparable to conditions observed in local background creeks and are unlikely to be attributable to residual Spill-related petroleum hydrocarbons.

**Riparian Birds and Mammals.** Potential risks to riparian birds and mammals due to residual exposures to aromatic and aliphatic fractions of TPH in Lower Red Butte Creek were identified. Note that exposures and risk estimates for aromatic and aliphatic fractions of TPH-diesel were comparable to or less than those measured/calculated for the local background creeks. Although statistical tests found no significant difference, visual examination of Q-Q plots suggest that TPH-motor oil concentrations appear to be greater in Red Butte Creek as compared to local background creeks. Risk estimates for the aliphatic fraction of TPH-motor oil<sup>35</sup> were greater than one and greater than risk estimates calculated for the local background creeks.

**Local Background Evaluation.** An evaluation of local background creeks found that concentrations of several PAHs and TPH-diesel in creek bed sediment of Lower Red Butte Creek were comparable or less than concentrations in creek bed sediment of local background creeks. Similarly, this evaluation found that concentrations of several PAHs and TPH-diesel in soil/sediment of Lower Red Butte Creek were comparable or less than concentrations in soil/sediments of local background creeks. Although visual examination of Q-Q plots found inconsistencies with the statistical analyses, an analysis of PAH composition suggests a source of PAHs that is consistent with urban runoff (McDaniel-Lambert 2012). This evaluation of the local background conditions suggest that petroleum hydrocarbons detected in Red Butte Creek may not be Spill-related.

**Uncertainties.** Uncertainties associated with the risk analyses were identified for this SLERA. To reduce uncertainties, focused verification of this SLERA may be considered. However, given potential risks to biota in Lower Red Butte Creek, for the most part, appear to be comparable to or less than risks for conditions in local background creeks, the need for verification may not be considered essential to support decision-making.

---

<sup>35</sup> The aromatic fraction of TPH-motor oil could not be quantitatively evaluated due to the lack of a mammalian TRV.



**10.0 LITURATURE CITED**

Agency for Toxic Substances and Disease Registry. 1999. *Toxicological profile for total petroleum hydrocarbons (TPH)*. Division of Toxicology/ Toxicology Information Branch, U.S. Department of Health and Human Services. Atlanta, GA.

Beyer, W.N., E.E. Connor, and S. Gerould. 1994. Estimates of Soil Ingestion by Wildlife. *Journal of Wildlife Management* 58(2):375-382.

Bio-West, Inc. 2010. Salt Lake City Riparian Corridor Study: Final Red Butte Creek Management Plan. Prepared for the Salt Lake City Department of Public Utilities. Salt Lake City, UT.

Calabrese, E.J., and L.A. Baldwin. 1993. *Performing Ecological Risk Assessments*. Lewis Publishers, Chelsea, MI.

California Environmental Protection Agency (Cal/EPA). 1996. Guidance for Ecological Risk Assessment at Hazardous Waste Sites and Permitted Facilities. California Department of Toxic Substances Control. July 4.

California Environmental Protection Agency (Cal/EPA). 2009. Evaluating Human Health Risks from Total Petroleum Hydrocarbons (TPH). Interim Guidance. Human and Ecological Risk Division. California Department of Toxic Substances Control. June 16.

Chevron Pipe Line (CPL). 2011. Red Butte Creek Crude Oil Spill Water, Sediment and Macroinvertebrate Sampling Plan. Version 17. Salt Lake City, Utah.

Donkin, P. 1994. Quantitative-Structure-Activity Relationships. In: *Handbook of Ecotoxicology* (P. Calow, ed.). Blackwell Scientific Publications. Oxford, England.

Douben, P.E.T. (ed.) 2003. *PAHs: An Ecotoxicological Perspective*. John Wiley & Sons Ltd. London, England.

ENTACT LLC (ENTACT). 2010. Removal Action Work Plan, Revision 2 for The Chevron Pipe Line Red Butte Release Mp 174.5, Salt Lake City, UT. Prepared for Chevron Environmental Management Company (CEMC).

Jones, D.S., G.W. Suter II, and R.N. Hull. 1997. Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Sediment-Associated Biota: 1997 revision. Health Sciences Research Division, Oak Ridge National Laboratory (ORNL). Oak Ridge, TN.

- MacDonald, D.D., C.G. Ingersoll, T.A. Berger. 2000. Development of Evaluation of Consensus-Based Sediment Quality Guidelines for Freshwater Ecosystems. *Archives of Environmental Contamination and Toxicology*. 39: 20-31.
- Massachusetts Department of Environmental Protection (MaDEP). 2002. Characterizing Risks Posed by Petroleum Contaminated Sites: Implementation of the MaDEP VPH/EPH Approach. Boston, MA.
- Massachusetts Department of Environmental Protection (MaDEP). 2007. Sediment Toxicity of Petroleum Hydrocarbon Fractions. Boston, MA.
- McDaniel-Lambert, Inc. (McDaniel-Lambert). 2012. Draft Human Health Risk Assessment, Red Butte Creek, Salt Lake City, Utah. Prepared for the Division of Water Resources, Utah Department of Environmental Quality.
- National Oceanic and Atmospheric Administration (NOAA). 2008. Sediment Quick Reference Tables (SQuiRT).
- Nirmalakhandan, N., and R.E. Speece. 1988. Structure-Activity Relationships. *Environ. Sci. Technol.* 22:606-615.
- Peterle, T.J. 1991. *Wildlife Toxicology*. Van Nostrand Reinhold, New York.
- Salt Lake County (SLCO). 2009. 2009 Salt Lake Countywide Water Quality Stewardship Plan. Salt Lake City, Salt Lake County, UT.
- Sample, B.E., D.M. Opresko, and G.W. Suter, II. 1996. Toxicological Benchmarks for Wildlife: 1996 revision. Health Sciences Research Division, Oak Ridge National Laboratory (ORNL). Oak Ridge, TN.
- Suter, G.W. II, and C.L. Tsao. 1996. Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Aquatic Biota: 1996 revision. Health Sciences Research Division, Oak Ridge National Laboratory (ORNL). Oak Ridge, TN.
- Total Petroleum Hydrocarbon Criteria Working Group (TPHCWG). 1997a. A risk-based approach for the management of total petroleum hydrocarbons in soil.
- Total Petroleum Hydrocarbon Criteria Working Group (TPHCWG). 1997b. Total petroleum hydrocarbon criteria working group series. Volume 4: Development of fraction specific reference doses (RfDs) and reference concentrations (RfCs) for total petroleum hydrocarbons (TPH). Amherst Scientific Publishers. Amherst, MA.

- United States Environmental Protection Agency (USEPA). 1989. Risk Assessment Guidance for Superfund (ERAGS). EPA/540/1-89/002. Office of Emergency and Remedial Response, United States Environmental Protection Agency. Washington, D.C.
- USEPA. 1991. Summary Report on Issues in Ecological Risk Assessment. EPA/625/3-91/018. Risk Assessment Forum, U.S. Environmental Protection Agency, Washington, D.C.
- USEPA. 1992a. Framework for Ecological Risk Assessment. EPA/630/R-92/001. Risk Assessment Forum, U.S. Environmental Protection Agency. Washington, D.C.
- USEPA. 1992b. Guidance for Data Usability in Risk Assessment. EPA-540/G-90-008. Office of Emergency and Remedial Response, U.S. Environmental Protection Agency. Washington, D.C.
- USEPA. 1993. Wildlife Exposure Factors Handbook. EPA/600/R-93/187. Office of Research and Development, U.S. Environmental Protection Agency. Washington, D.C.
- USEPA. 1997. Ecological Risk Assessment Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessments. Interim Final. EPA-540-R-97-006. Solid Waste and Emergency Response, U.S. Environmental Protection Agency, Washington, DC.
- USEPA. 1998. Guidelines for Ecological Risk Assessment. EPA/630/R-95/002F. Risk Assessment Forum, U.S. Environmental Protection Agency. Washington, D.C.
- USEPA. 1999a. National Functional Guidelines for Organic Data Review. USEPA 540/R-99-008. OSWER 9240.1-05A-P. October.
- USEPA. 1999b. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish. Second Edition. EPA/841/B-99/002. Office of Office of Wetlands, Oceans, and Watersheds, U.S. Environmental Protection Agency. Washington, D.C.
- USEPA. 2001a. Risk Assessment Guidance for Superfund: Volume I Human Health Evaluation Manual – Part D, Standardized Planning, Reporting, and Review of Superfund Risk Assessments. Office of Emergency and Remedial Response, Washington, DC. Publication 9285.7-47. December.

USEPA. 2002. Guidance for Comparing Background and Chemical Concentrations in Soil for CERCLA Sites. EPA 540-R-01-003. Office of Solid Waste and Emergency Response, U.S. Environmental Protection Agency, Washington, D.C.

USEPA. 2004. National Functional Guidelines for Inorganic Data Review. USEPA 540-R-04-004. OSWER 9240.1-45. October.

USEPA. 2005. Guidance for Developing Ecological Soil Screening Levels. Office of Solid Waste and Emergency Response. Directive 9285.7-55. February 2005. Updated 2007.

USEPA (Singh, A., Maichle, R.W. and S. Lee). 2006. On the Computation of a 95% Upper Confidence Limit of the Unknown Population Mean Based Upon Data Sets with Below Detection Limit Observations. EPA/600/R-06/022.

USEPA. 2007. Ecological Soil Screening Levels for Polycyclic Aromatic Hydrocarbons (PAHs). Interim Final. Office of Solid Waste and Emergency Response, Washington, D.C.

USEPA. 2009. National Recommended Water Quality Criteria. Office of Water, Office of Science and Technology (4304T).  
<http://water.epa.gov/scitech/swguidance/standards/current/>


USEPA (Singh, A., and A.K. Singh). 2010. ProUCL Version 4.0 Technical Guide. EPA/600/R-07/04. Office of Research and Development, U.S. Environmental Protection Agency, Las Vegas, NV.



Utah Administrative Code. Rule R315-101-5, Health Evaluation Criteria, Risk Assessment.

Utah Department of Environmental Quality (UDEQ). 2005. TPH Fractionation at Leaking Underground Storage Tank Sites, Final Draft. Updated November 2011.


## FIGURES

• This page intentionally left blank •















e Creek



Date: 1/05/12



metrics



Date:  
11/05/12





• This page intentionally left blank •

## TABLES

(Tables 8-1a,b through 8-6a,b)

• This page intentionally left blank •



*Table 8-1a  
Risk Calculations for Aquatic Biota in Red Butte Creek  
Ecological Risk Assessment  
Red Butte Creek  
Salt Lake City, Utah*

Water COPEC	Media EPC	TRV (ug/L)	Hazard <sup>1</sup> Quotient
	Water (ug/L)		
None			

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration. The maximum concentration was used as the EPC for this Screening Level evaluation.

BAF - Bioaccumulation Factor

NA - Not applicable or unavailable

TRV - Toxicity Reference Value

HQ - Hazard Quotient

1 -Hazard Quotient = Total Dose/TRV

*Table 8-1b  
 Risk Calculations for Aquatic Biota in Local Background Creeks  
 Ecological Risk Assessment  
 Red Butte Creek  
 Salt Lake City, Utah*

Water COPEC	Media EPC	TRV (ug/L)	Hazard <sup>1</sup> Quotient
	Water (ug/L)		
None			

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration. The maximum concentration was used as the EPC for this Screening Level evaluation.

NA - Not applicable or unavailable

TRV - Toxicity Reference Value

HQ - Hazard Quotient

1 -Hazard Quotient = Total Dose/TRV

Table 8-2a  
 Risk Calculations for Benthic Invertebrates in Red Butte Creek  
 Ecological Risk Assessment  
 Red Butte Creek  
 Salt Lake City, Utah

Sediment COPEC	Media EPC	Sediment TRV		Hazard Quotient <sup>1</sup>	
	Sediment (mg/kg)	TEC (mg/kg)	PEC (mg/kg)	TEC	PEC
Low Molecular Weight PAHs					
Anthracene	0.090	0.057	0.85	1.6	0.11
High Molecular Weight PAHs					
Benzo(a)anthracene	0.063	0.11	1.1	0.58	0.060
Benzo(a)pyrene	0.085	0.15	1.5	0.57	0.059
Benzo(b)fluoranthene	0.082	0.027	1.1	3.0	0.078
Benzo(g,h,i)perylene	0.074	0.17	1.1	0.44	0.071
Benzo(k)fluoranthene	0.044	0.027	1.1	1.6	0.041
Dibenzo(a,h)anthracene	0.058	0.033	1.1	1.8	0.055
Indeno(1,2,3-cd)pyrene	0.12	0.017	1.1	7.0	0.12
Pyrene	0.087	0.20	1.5	0.45	0.058
TPH Diesel					
Aromatics	49	0.29	--	172	--
Aliphatics	49	9.9	--	5.0	--
TPH Motor Oil					
Aromatics	53	--	--	--	--
Aliphatics	53	31	--	1.7	--
		Low Molecular Weight PAHs HI =		1.6	0.11
		High Molecular Weight PAHs HI =		15	0.54
		TPH HI =		179	--

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration.

NA - Not applicable or unavailable

HQ - Hazard Quotient

<sup>1</sup> -Hazard Quotient = Total Dose/TRV

Table 8-2b  
 Risk Calculations for Benthic Invertebrates in Local Background Creeks  
 Ecological Risk Assessment  
 Red Butte Creek  
 Salt Lake City, Utah

Sediment COPEC	Media EPC	Sediment TRV		Hazard Quotient <sup>1</sup>	
	Sediment (mg/kg)	TEC (mg/kg)	PEC (mg/kg)	TEC	PEC
Low Molecular Weight PAHs					
Anthracene	0.0059	0.057	0.85	0.10	0.0070
High Molecular Weight PAHs					
Benzo(a)anthracene	0.056	0.11	1.1	0.52	0.054
Benzo(a)pyrene	0.040	0.15	1.5	0.27	0.028
Benzo(b)fluoranthene	0.057	0.027	1.1	2.1	0.054
Benzo(g,h,i)perylene	0.031	0.17	1.1	0.18	0.030
Benzo(k)fluoranthene	0.027	0.027	1.1	0.99	0.026
Dibenzo(a,h)anthracene	--	0.033	1.1	--	
Indeno(1,2,3-cd)pyrene	0.033	0.017	1.1	1.9	0.031
Pyrene	0.11	0.20	1.5	0.55	0.072
TPH Diesel	105				
Aromatics	53	0.29	--	184	--
Aliphatics	53	9.9	--	5.3	--
TPH Motor Oil	67				
Aromatics	34	--	--	--	--
Aliphatics	34	31	--	1.1	--
		Low Molecular Weight PAHs HI =		0.10	0.01
		High Molecular Weight PAHs HI =		6.5	0.29
		TPH HI =		190	--

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration.

NA - Not applicable or unavailable

HQ - Hazard Quotient

1 -Hazard Quotient = Total Dose/TRV

*Table 8-3a  
Risk Calculations for Mallard at Red Butte Creek  
Ecological Risk Assessment  
Red Butte Creek  
Salt Lake City, Utah*

Sediment COPEC	Media EPC		Bioaccumulation Model <sup>1</sup>	Food EPC	Dose <sup>2</sup>			Total Dose (mg/kg-day)	Avian TRV (mg/kg-day)	Hazard <sup>3</sup> Quotient
	Soil (mg/kg)	Surface Water (mg/L)	BAF	Plant (mg/kg)	Incidental Sediment Ingestion (mg/kg-day)	Surface Water Ingestion (mg/kg-day)	Plant Ingestion (mg/kg-day)			
TPH-Diesel										
Aromatics	51	NA	1.2	62	0.095	NA	3.5	3.6	NA	--
Aliphatics	51	NA	0.54	27	0.095	NA	1.5	1.6	NA	--
TPH Motor Oil										
Aromatics	48	NA	1.2	58	0.089	NA	3.3	3.4	NA	--
Aliphatics	48	NA	0.54	26	0.089	NA	1.4	1.5	NA	--

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration. The maximum concentration was used as the EPC for this Screening Level evaluation.

BAF - Bioaccumulation Factor

NA - Not applicable or unavailable

TRV - Toxicity Reference Value

HQ - Hazard Quotient

1 -Food EPCs calculated using a point estimate bioaccumulation factor.

2 -See text for dose calculations.

3 -Hazard Quotient = Total Dose/TRV

*Table 8-3b  
Risk Calculations for Mallard in Local Background Creeks  
Ecological Risk Assessment  
Red Butte Creek  
Salt Lake City, Utah*

Sediment COPEC	Media EPC		Bioaccumulation Model <sup>1</sup>	Food EPC	Dose <sup>2</sup>			Total	Avian	Hazard <sup>3</sup> Quotient
	Soil (mg/kg)	Surface Water (mg/L)	BAF	Plant (mg/kg)	Incidental Sediment Ingestion (mg/kg-day)	Surface Water Ingestion (mg/kg-day)	Plant Ingestion (mg/kg-day)	Dose (mg/kg-day)	TRV (mg/kg-day)	
TPH Diesel	154									
Aromatics	77	NA	1.2	94	0.142	NA	5.2	5.4	NA	--
Aliphatics	77	NA	0.54	41	0.142	NA	2.3	2.5	NA	--
TPH Motor Oil	98									
Aromatics	49	NA	1.2	60	0.091	NA	3.3	3.4	NA	--
Aliphatics	49	NA	0.54	26	0.091	NA	1.5	1.6	NA	--

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration. The maximum concentration was used as the EPC for this Screening Level evaluation.

BAF - Bioaccumulation Factor

NA - Not applicable or unavailable

TRV - Toxicity Reference Value

HQ - Hazard Quotient

1 -Food EPCs calculated using a point estimate bioaccumulation factor.

2 -See text for dose calculations.

3 -Hazard Quotient = Total Dose/TRV

*Table 8-4a  
Risk Calculations for Sandpiper at Red Butte Creek  
Ecological Risk Assessment  
Red Butte Creek  
Salt Lake City, Utah*

Sediment COPEC	Media EPC		Bioaccumulation Model <sup>1</sup>	Food EPC	Dose <sup>2</sup>			Total	Avian	Hazard <sup>3</sup> Quotient
	Sediment (mg/kg)	Surface Water (mg/L)	BAF	Invert (mg/kg)	Incidental Sediment Ingestion (mg/kg-day)	Surface Water Ingestion (mg/kg-day)	Invertebrate Ingestion (mg/kg-day)	Dose (mg/kg-day)	TRV (mg/kg-day)	
TPH-Diesel										
Aromatics	51	NA	1431	73192	0.68	NA	11930	11931	NA	--
Aliphatics	51	NA	17	883	0.68	NA	144	145	NA	--
TPH Motor Oil										
Aromatics	48	NA	1431	68592	0.64	NA	11180	11181	NA	--
Aliphatics	48	NA	17	827	0.64	NA	135	136	NA	--

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration. The maximum concentration was used as the EPC for this Screening Level evaluation.

BAF - Bioaccumulation Factor

NA - Not applicable or unavailable

TRV - Toxicity Reference Value

HQ - Hazard Quotient

1 -Food EPCs calculated using a point estimate bioaccumulation factor.

2 -See text for dose calculations.

3 -Hazard Quotient = Total Dose/TRV

*Table 8-4b  
Risk Calculations for Sandpiper in Local Background Creeks  
Ecological Risk Assessment  
Red Butte Creek  
Salt Lake City, Utah*

Sediment COPEC	Media EPC		Bioaccumulation Model <sup>1</sup>	Food EPC	Dose <sup>2</sup>			Total	Avian	Hazard <sup>3</sup> Quotient
	Sediment (mg/kg)	Surface Water (mg/L)	BAF	Invert (mg/kg)	Incidental Sediment Ingestion (mg/kg-day)	Surface Water Ingestion (mg/kg-day)	Invertebrate Ingestion (mg/kg-day)	Dose (mg/kg-day)	TRV (mg/kg-day)	
TPH Diesel	154									
Aromatics	77	NA	1431	110182	1.03	NA	17960	17961	NA	--
Aliphatics	77	NA	17	1329	1.03	NA	217	218	NA	--
TPH Motor Oil	98									
Aromatics	49	NA	1431	70338	0.66	NA	11465	11466	NA	--
Aliphatics	49	NA	17	848	0.66	NA	138	139	NA	--

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration. The maximum concentration was used as the EPC for this Screening Level evaluation.

BAF - Bioaccumulation Factor

NA - Not applicable or unavailable

TRV - Toxicity Reference Value

HQ - Hazard Quotient

1 -Food EPCs calculated using a point estimate bioaccumulation factor.

2 -See text for dose calculations.

3 -Hazard Quotient = Total Dose/TRV



*Table 8-5a  
Risk Calculations for Muskrat at Red Butte Creek  
Ecological Risk Assessment  
Red Butte Creek  
Salt Lake City, Utah*

Soil COPEC	Media EPC		Bioaccumulation Model <sup>1</sup>	Food EPC	Dose <sup>2</sup>			Total	Mammal	Hazard <sup>3</sup> Quotient
	Soil (mg/kg)	Surface Water (mg/L)	BAF	Plant (mg/kg)	Incidental Soil Ingestion (mg/kg-day)	Surface Water Ingestion (mg/kg-day)	Plant Ingestion (mg/kg-day)	Dose (mg/kg-day)	TRV (mg/kg-day)	
TPH-Diesel										
Aromatics	51	NA	1.22	62	1.4	NA	19	20	3.0	6.7
Aliphatics	51	NA	0.54	27	1.4	NA	8.2	9.7	10	0.97
TPH Motor Oil										
Aromatics	48	NA	1.2	58	1.4	NA	17	19	--	--
Aliphatics	48	NA	0.54	26	1.4	NA	7.7	9.1	60	0.15
									TPH HI =	7.8

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration. The maximum concentration was used as the EPC for this Screening Level evaluation.

BAF - Bioaccumulation Factor

NA - Not applicable or unavailable

TRV - Toxicity Reference Value

HQ - Hazard Quotient

1 -Food EPCs calculated using a point estimate bioaccumulation factor.

2 -See text for dose calculations.

3 -Hazard Quotient = Total Dose/TRV

*Table 8-5b  
Risk Calculations for Muskrat in Local Background Creeks  
Ecological Risk Assessment  
Red Butte Creek  
Salt Lake City, Utah*

Soil COPEC	Media EPC		Bioaccumulation Model <sup>1</sup>	Food EPC	Dose <sup>2</sup>			Total	Mammal	Hazard <sup>3</sup> Quotient
	Soil (mg/kg)	Surface Water (mg/L)	BAF	Plant (mg/kg)	Incidental Soil Ingestion (mg/kg-day)	Surface Water Ingestion (mg/kg-day)	Plant Ingestion (mg/kg-day)	Dose (mg/kg-day)	TRV (mg/kg-day)	
TPH Diesel	154									
Aromatics	77	NA	1.22	94	2.2	NA	28	30	3.0	10
Aliphatics	77	NA	0.54	41	2.2	NA	12.4	15	10	1.5
TPH Motor Oil	98									
Aromatics	49	NA	1.2	60	1.39	NA	18	19	--	--
Aliphatics	49	NA	0.54	26	1.39	NA	7.9	9.3	60	0.16
									TPH HI =	12

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration. The maximum concentration was used as the EPC for this Screening Level evaluation.

BAF - Bioaccumulation Factor

NA - Not applicable or unavailable

TRV - Toxicity Reference Value

HQ - Hazard Quotient

1 -Food EPCs calculated using a point estimate bioaccumulation factor.

2 -See text for dose calculations.

3 -Hazard Quotient = Total Dose/TRV

*Table 8-6a  
Risk Calculations for Raccoon at Red Butte Creek  
Ecological Risk Assessment  
Red Butte Creek  
Salt Lake City, Utah*

Sediment COPEC	Media EPC		Bioaccumulation Model <sup>1</sup>	Food EPC	Dose <sup>2</sup>			Total Dose (mg/kg-day)	Mammal TRV (mg/kg-day)	Hazard <sup>3</sup> Quotient
	Sediment (mg/kg)	Surface Water (mg/L)	BAF	Inverts (mg/kg)	Incidental Sediment Ingestion (mg/kg-day)	Surface Water Ingestion (mg/kg-day)	Invertebrate Ingestion (mg/kg-day)			
TPH-Diesel										
Aromatics	51	NA	1431	73192	2.6	NA	39304	39307	3.0	13102
Aliphatics	51	NA	17	883	2.6	NA	474	477	10	48
TPH Motor Oil										
Aromatics	48	NA	1431	68592	2.4	NA	36834	36836	--	--
Aliphatics	48	NA	17	827	2.4	NA	444	447	60	7.4
									TPH HI =	13157

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration. The maximum concentration was used as the EPC for this Screening Level evaluation.

BAF - Bioaccumulation Factor

NA - Not applicable or unavailable

TRV - Toxicity Reference Value

HQ - Hazard Quotient

1 -Food EPCs calculated using a point estimate bioaccumulation factor.

2 -See text for dose calculations.

3 -Hazard Quotient = Total Dose/TRV

*Table 8-6b  
Risk Calculations for Raccoon in Local Background Creeks  
Ecological Risk Assessment  
Red Butte Creek  
Salt Lake City, Utah*

Sediment COPEC	Media EPC		Bioaccumulation Model <sup>1</sup>	Food EPC	Dose <sup>2</sup>			Total Dose (mg/kg-day)	Mammal TRV (mg/kg-day)	Hazard <sup>3</sup> Quotient
	Sediment (mg/kg)	Surface Water (mg/L)	BAF	Inverts (mg/kg)	Incidental Sediment Ingestion (mg/kg-day)	Surface Water Ingestion (mg/kg-day)	Invertebrate Ingestion (mg/kg-day)			
TPH Diesel	154									
Aromatics	77	NA	1431	110182	3.9	NA	59168	59172	3.0	19724
Aliphatics	77	NA	17	1329	3.9	NA	714	718	10	72
TPH Motor Oil	98									
Aromatics	49	NA	1431	70338	2.5	NA	37771	37774	--	--
Aliphatics	49	NA	17	848	2.5	NA	456	458	60	7.6
									TPH HI =	19803

COPEC - Chemical of Potential Ecological Concern

EPC - Exposure Point Concentration. The maximum concentration was used as the EPC for this Screening Level evaluation.

BAF - Bioaccumulation Factor

NA - Not applicable or unavailable

TRV - Toxicity Reference Value

HQ - Hazard Quotient

1 -Food EPCs calculated using a point estimate bioaccumulation factor.

2 -See text for dose calculations.

3 -Hazard Quotient = Total Dose/TRV