#### **Chapter 2.9.1 Cedar/Beaver Watershed Management Unit Assessment**

#### 2.9.1. Introduction

The Cedar / Beaver Watershed Management Unit includes all streams located in the U.S.G.S Hydrological Units (HUCs) listed in Table 2.9.1. There are not many streams within this unit with the major streams being the Beaver River, Coal Creek, Shoal Creek and Pinto Creek.

| Table 2.9.1. U.S.G.S. Hydrological Units in the Cedar/Beaver<br>Watershed Management Unit. |                             |  |  |
|--|-----------------------------|--|--|
| Hydrological Unit Code Hydrological Unit Name  |                             |  |  |
| 16030006   | Escalante Desert            |  |  |
| 16030007   | Beaver Bottoms-Upper Beaver |  |  |
| 16030008   | Lower Beaver                |  |  |

#### 2.9.2. Water Quality Assessment Results

**2.9.2.1. Overall Beneficial Use Support**—Data collected between January 1, 2002 and December 31, 2006, including the intensive survey were used to determine beneficial use support. Benthic macroinvertebrate data were used for the first time in making beneficial use assessments (Chapter 2.15).

Beneficial use support assessments are made by comparing data against numeric standards established for each beneficial use. Figure 2.9.2 is a map of the designated beneficial uses assigned to the stream and river Assessment Units. Assessments using benthic macroinvertebrate data are based upon the State's narrative standard.

An assessment of support for at least one beneficial use was made for 289.2 stream miles. Of those assessed, 231.6 miles (80.1%) are fully supporting, and all the beneficial uses

Full 80.1% Not 289 miles assessed

Overall Benficial Use Support

Figure 2.9.1. Overall beneficial use support.

assessed and (19.9%) are not supporting at least one designated beneficial use. The overall beneficial use assessment is shown in Figure 2.9.1.

**2.9.2.2. Beneficial Use Assessment By Categories--** The number of stream miles assessed by categories is listed in Table 2.9.2. Figure 2.9.3 is a map of the assessment categories that rivers and streams were assigned to after the beneficial uses were evaluated. An Assessment Unit (AU) can be in more than one category.

| Table. 2.9.2.Stream Miles By Assessment Category – Cedar/Beaver |  |              |  |
|---|--|--------------|--|
| Category  | Category Definition  | Stream Miles |  |
| 1   | All beneficial uses fully supported.   | 0            |  |
| 2   | Beneficial uses assessed are fully supported.  | 231.61       |  |
| 3A  | No data or insufficient data to make an assessment.  | 51.2         |  |
| 3B  | Lakes that are not supported for one cycle only.   | 0.0          |  |
| 3C  | Insufficient data to assess but an assessment plan is in place.                                      | 0.0          |  |
| 4A  | Approved TMDL  | 57.7         |  |
| 4B  | Pollution control requirements are expected to result in full beneficial use support in near future. | 0            |  |
| 4C  | Impaired by pollution, no TMDL required.   | 57.7         |  |
| 5   | Impaired by pollutant, TMDL required.  | 0.0          |  |

**2.9.2.3. Individual Beneficial Use Support--** Individual beneficial use support is listed in Table 2.9.3. For aquatic life use support, 289.2 miles (80.1%) are fully supporting and 57.6 miles (19.9%) are not supporting this beneficial use. Of the 275.7 stream miles assessed for agricultural use, 218.1 (79.1%) were assessed as fully supporting and 57.6 miles as (20.9%) not supporting this designated beneficial use. The 57.6 miles assessed for swimming and secondary contact are not supporting this beneficial use because of pH.

| Table 2.9.3 Individual Beneficial Use Support – Cedar/Beaver Watershed |          |            |            |        |  |  |
|--|----------|------------|------------|--------|--|--|
| Management Unit (Stream Miles) Classification - 2008                   |          |            |            |        |  |  |
|  |          |            |            |        |  |  |
|  | Size     | Size Fully | Size Not   |        |  |  |
|  | Assessed | Supporting | Supporting | Totals |  |  |
| Use  |          |            |            |        |  |  |
| Aquatic Life   | 289.2    | 231.6      | 57.6       | 317.3  |  |  |
| Fish Consumption   | 0.0      | 0.0        | 0.0        | 0.0    |  |  |
| Swimming   | 57.6     | 0.0        | 57.6       | 57.6   |  |  |
| Secondary Contact  | 57.6     | 0.0        | 57.6       | 57.6   |  |  |
| Drinking Water   | 0.0      | 0.0        | 0.0        | 0.0    |  |  |
| Agricultural   | 275.7    | 218.1      | 57.6       | 275.7  |  |  |
|  |          |            |            |        |  |  |
| Use  |          |            |            |        |  |  |
| Aquatic Life   |          | 80.1%      | 19.9%      | 100.0% |  |  |
| Fish Consumption   |          | 0.0%       | 0.0%       | 0.0%   |  |  |
| Swimming   |          | 0.0%       | 100.0%     | 100.0% |  |  |
| Secondary Contact  |          | 0.0%       | 100.0%     | 100.0% |  |  |
| Drinking Water   |          | 0.0%       | 0.0%       | 0.0%   |  |  |
| Agricultural   |          | 80.1%      | 19.9%      | 100.0% |  |  |




Figure 2.9.2. River and stream designated beneficial use classes – Cedar/Beaver Watershed Management Unit.




Figure 2.9.3. Beneficial use assessment by category - Cedar / Beaver Watershed Management Unit.

**2.9.2.4. Total Waters Impaired by Various Causes**— The causes of impairment are listed in Table 2.9.4. The causes of impairment are nutrients (total phosphorus), thermal modification, pH and habitat alterations. The percent of miles impacted by various causes is illustrated in Figure 2.9.4. The relative impact of these causes is shown in Figure 2.9.5.

**2.9.2.5. Total Waters Impaired by Various Sources**—The number of stream miles impacted by sources are listed in table 2.9.5. The sources of impairment are agricultural activities, hydromodification, habitat modification, and unknown sources as shown in Figure 2.9.6. The relative percent impairment by sources is illustrated in Figure 2.9.7.

The impaired Assessment Units are listed in Table 2.9.6. The table includes the class of beneficial uses impaired and the pollutant or pollution causing the impairment.

This page left blank on purpose.

| Table 2.9.4. Total Waters Impaired by Various Cause<br>Categories – Cedar/Beaver Watershed<br>Management Unit. |              |  |
|--|--------------|--|
| Cause Category   | Stream Miles |  |
| Cause unknown  | 0.0          |  |
| Unknown toxicity   | 0.0          |  |
| Pesticides   | 0.0          |  |
| Priority organics  | 0.0          |  |
| Nonpriority organics   | 0.0          |  |
| Metals   | 0.0          |  |
| Ammonia  | 0.0          |  |
| Chlorine   | 0.0          |  |
| Other inorganics   | 0.0          |  |
| Nutrients  | 57.7         |  |
| рН   | 57.7         |  |
| Siltation/Sediments  | 0.0          |  |
| Organic enrichment/low DO  | 0.0          |  |
| Salinity/TDS/Chlorides   | 0.0          |  |
| Thermal modifications  | 57.7         |  |
| Flow alterations   | 0.0          |  |
| Other habitat alterations  | 57.7         |  |
| Pathogen Indicators  | 0.0          |  |
| Radiation  | 0.0          |  |
| Oil and grease   | 0.0          |  |
| Taste and odor   | 0.0          |  |
| Noxious aquatic plants   | 0.0          |  |
| Total Toxics   | 0.0          |  |
| Turbidity  | 0.0          |  |
| Benthic Macroinvertebrates   | 0.0          |  |
| Other (Specify)  | 0.0          |  |

| Table 2.9.5. Total Waters Impaired by Various Source<br>Categories – Cedar/Beaver Watershed<br>Management Unit. |      |  |  |
|---|------|--|--|
|   |      |  |  |
| Industrial Point Sources  | 0.0  |  |  |
| Municipal Point Sources   | 0.0  |  |  |
| Combined Sewer Overflow   | 0.0  |  |  |
| Agriculture   | 57.7 |  |  |
| Silviculture  | 0.0  |  |  |
| Construction  | 0.0  |  |  |
| Urban Runoff/Storm Sewers   | 0.0  |  |  |
| Resource Extraction   | 0.0  |  |  |
| Land Disposal   | 0.0  |  |  |
| Hydromodification   | 57.7 |  |  |
| Habitat Modification  | 57.7 |  |  |
| Marinas   | 0.0  |  |  |
| Atmospheric Deposition  | 0.0  |  |  |
| Contaminated Sediments  | 0.0  |  |  |
| Unknown Source  | 0.0  |  |  |
| Natural Sources   | 57.7 |  |  |
| Reservoir Releases  | 0.0  |  |  |
| Recreation  | 0.0  |  |  |
| Aquaculture   | 0.0  |  |  |
| Extreme Drought   | 0.0  |  |  |

## Percent of Stream Miles Affected By Causes

2008 Integrated Report Assessment - Cedar / Bearver Management Unit




Figure 2.9.4. Percent of assessed stream miles impacted by various causes – Cedar/Beaver Watershed Management Unit.

# Causes of Stream Water Quality Impairments 2008 Integrated Report Assessment - Cedar / Beaver Mangement Unit




Figure 2.9.5. Relative percent impact by causes on water quality – Cedar/Beaver Watershed Management Unit.

## Percent of Stream Miles Affected By Sources

2008 Integrated Report Assessement - Cedar / Beaver Watershed Management Unit




Figure 2.9.6. Percent of assessed stream miles impacted by various sources – Cedar/Beaver Watershed Management Unit.

# Sources of Stream Water Quality Impairment

2008 Integrated Report Assessment - Cedar / Beaver Watershed Management Unit




Figure 2.9.7. Relative percent contribution of causes on stream water quality – Cedar/Beaver Watershed Management Unit.

|                | Table 2.9.6. Impaired Waters Located in the Cedar/Beaver Watershed Management Unit. |   |            |          |            |                           |                 |
|----------------|---|---|------------|----------|------------|---------------------------|-----------------|
|                |   |   | Beneficial |          | Beneficial |                           |                 |
| Assessment     | Assessment  | Assessment  | Use        |          | Use        | Pollutant                 |                 |
| Unit           | Unit  | Unit  | Class      | Support  | Assessment | Or                        | Stream          |
| ID             | Name  | Description   | Impaired   | Category | Category   | Pollution                 | Stream<br>Miles |
| UT16030007-002 | Beaver River-2  | Beaver River and tributaries from<br>Minersville Reservoir to USFS boundary | 2B, 3A, 4  | NS       | 4A         | pH                        | 57.57           |
| UT16030007-002 | Beaver River-2  | Beaver River and tributaries from<br>Minersville Reservoir to USFS boundary | 3A         | NS       | 4A         | Total Phosphorus          | 57.57           |
| UT16030007-002 | Beaver River-2  | Beaver River and tributaries from<br>Minersville Reservoir to USFS boundary | 3A         | NS       | 4A         | Thermal Modifications     | 57.57           |
| UT16030007-002 | Beaver River-2  | Beaver River and tributaries from<br>Minersville Reservoir to USFS boundary | 3A         | NS       | 4C         | Other Habitat Alterations | 57.57           |